herpy

ESCINCOS

Southern White's Skink

Liopholis whitii

A medium-sized, diurnal rock skink from cool temperate southern Australia and Tasmania. *Liopholis whitii* lives in rock crevices and burrows under stones, lives in small family groups and gives birth to live young. It needs an enclosure with many narrow hiding places, a warm basking spot, a cool night and patience when putting animals together.

Familia: Scincidae Nombre anterior: *Egernia whitii; Scincus whitii*

DATOS CLAVE

EL ANIMAL

Tamaño adulto	**8-10 cm** longitud hocico-cloaca
Esperanza de vida	**10-15 años**
Activo	**De día**
Modo de vida	**Rupícola y excavador**
Hábitat	**Terreno rocoso y bosque y prado templados**
Dificultad	**Intermedio**

CUIDADOS

Terrario mín.	**90 × 45 × 45 cm** L × An × Al para 2 adultos
Temperatura, día	**20-26 °C**
Punto caliente	**30-35 °C**
Temperatura, noche	**12-18 °C**
Humedad	**40-60 %**
Agua	**Bebedero**
UVB	**Recomendado** Zona Ferguson 2
Alimentación	**Insectos, fruta y plantas**
Se mantiene	**En pareja o en grupo**

CRÍA

Reproducción	**Vivípara**
Madurez sexual	**24-36 meses**
Tamaño de la camada	**1-4 crías**
Puestas por año	**1**
Gestación	**100-140 días**

PROTECCIÓN Y TRÁMITES

CITES	**Apéndice III**
Anexo de la UE	**Anexo C**
UICN	**Preocupación menor (LC)**

Cuidados de un vistazo

EL TERRARIO	Little is known about keeping *Liopholis whitii*. A single animal can be kept in 60 × 30 × 40 cm.
TEMPERATURA	By day the air should be 20-26 °C, and under a lamp at one end a flat stone should reach 30-35 °C measured with an infrared thermometer. At night the temperature should drop to 12-18 °C.
HUMEDAD Y AGUA	Humidity should be 40-60 %. Keep the substrate dry, but make part of it slightly damp under a stone or a piece of bark so the animals can choose. Do not mist daily.
ILUMINACIÓN Y UVB	The species is diurnal and basks on rocks. A UVB tube in a reflector above the basking spot should provide UVB corresponding to Ferguson zone 2.
ALIMENTACIÓN	*Liopholis whitii* is omnivorous. Offer insects such as mealworms, crickets and cockroaches, and supplement with finely chopped herbs and a little fruit or berries.
COMPORTAMIENTO Y MANIPULACIÓN	The species is territorial towards strangers and in the wild lives in small family groups whose members know each other.
ÁREA DE DISTRIBUCIÓN NATURAL	Australia.