Niceforo's Poison Frog

Ameerega ingeri

Ameerega ingeri is a small, diurnal poison dart frog of 20-27 mm from the foothills of the Andes in southern Colombia. The back is uniformly dark, the flanks have dark blue spots, and there are bright orange spots in the armpits and on the calves. The species is hardly ever kept, so care is based on the climate of its range and on related species.

Familia: Dendrobatidae Nombre anterior: *Dendrobates ingeri; Epipedobates ingeri; Phyllobates ingeri*

DATOS CLAVE

EL ANIMAL

Tamaño adulto	**20-27 mm** longitud hocico-cloaca
Esperanza de vida	**10-15 años**
Activo	**De día**
Modo de vida	**Terrestre**
Hábitat	**Selva tropical**
Toxinas cutáneas en la naturaleza	**Leves**
Toxinas cutáneas en cautividad	**Ninguna** Animales criados en cautividad. Los animales capturados en la naturaleza pueden conservar sus toxinas durante mucho tiempo
Dificultad	**Intermedio**

CRÍA

Reproducción	**Ovípara**
Desarrollo	**Renacuajos transportados al agua**
Lugar de puesta	**En el suelo**
Huevos por puesta	**4-16**
Eclosión	**10-22 días**
Periodo larvario	**70-90 días** desde la eclosión hasta la metamorfosis
Alimentación de las larvas	**Algas/plantas y alimento completo**
Cuidado parental	**Transporta los renacuajos**

CUIDADOS

Terrario mín.	**60 × 40 × 40 cm** L × An × Al para 2 adultos
Temperatura, día	**24-26 °C**
Temperatura, noche	**20-22 °C**
Humedad	**70-100 %**
Agua	**Bebedero** profundidad 1-2 cm
UVB	**Opcional** Zona Ferguson 1
Alimentación	**Pequeños invertebrados (colémbolos, moscas de la fruta)**
Se mantiene	**En pareja**

PROTECCIÓN Y TRÁMITES

CITES	**Apéndice II**
UICN	**Datos insuficientes (DD)**

Cuidados de un vistazo

EL TERRARIO	Almost nothing is known about the care of *Ameerega ingeri*, and the species is hardly ever kept.
TEMPERATURA	Keep the enclosure at 24-26 °C by day, and let the temperature fall to 20-22 °C at night. The temperature must not exceed 28 °C, because frogs tolerate heat poorly.
HUMEDAD Y PULVERIZACIÓN	Humidity should be 70-80 % in the middle of the day and close to 100 % in the morning and evening. A rain system or a fogger with a timer is easiest; otherwise mist two to three times a day with reverse-osmosis water or rainwater.
AGUA Y CALIDAD DEL AGUA	Provide a shallow water dish with 1-2 cm of water, so that the frogs can easily climb out of it, and with stones or moss at the edge. Use reverse-osmosis water, rainwater or dechlorinated tap water, and change it often.
ILUMINACIÓN Y UVB	The lighting is primarily for the plants: 10-12 hours a day with LED lighting with a plant-suitable spectrum. It is not documented that *Ameerega ingeri* needs UVB.
ALIMENTACIÓN	In the wild *Ameerega ingeri* lives mostly on small ants and other small arthropods from the leaf litter.
COMPORTAMIENTO Y MANIPULACIÓN	*Ameerega ingeri* is diurnal and most active in the morning. It lives on the ground in the leaf litter, but likes to sleep on leaves some way above the ground, so plants with broad leaves are an advantage.
ÁREA DE DISTRIBUCIÓN NATURAL	Colombia.