herpy

RANAS

Cowan's Mantella

Mantella cowanii

Mantella cowanii is a small, diurnal frog from the highlands of central Madagascar, glossy black with bright red to orange limbs. It grows to about 2-3 cm and is one of the most heat-sensitive in the genus: the enclosure must be kept cool all year round. The species is threatened in the wild, so only captive-bred animals should be considered.

Familia: Mantellidae

DATOS CLAVE

EL ANIMAL

Tamaño adulto	**2-3 cm** longitud hocico-cloaca
Esperanza de vida	**5-10 años**
Activo	**De día**
Modo de vida	**Terrestre**
Hábitat	**Bosque de montaña y humedal**
Toxinas cutáneas en la naturaleza	**Leves**
Toxinas cutáneas en cautividad	**Ninguna** Animales criados en cautividad. Los animales capturados en la naturaleza pueden conservar sus toxinas durante mucho tiempo
Dificultad	**Avanzado**

CRÍA

Reproducción	**Ovípara**
Desarrollo	**Renacuajos en el agua**
Lugar de puesta	**En el suelo**
Madurez sexual	**12-18 meses**
Huevos por puesta	**50-130**
Eclosión	**3-12 días**
Periodo larvario	**56-98 días** desde la eclosión hasta la metamorfosis
Alimentación de las larvas	**Algas/plantas y alimento completo**
Cuidado parental	**Ninguno**

CUIDADOS

Terrario mín.	**45 × 45 × 30 cm** L × An × Al para 4 adultos
Temperatura, día	**18-21 °C**
Temperatura, noche	**16-18 °C**
Humedad	**60-80 %**
Agua	**Bebedero**
UVB	**Opcional** Zona Ferguson 1
Alimentación	**Pequeños invertebrados (colémbolos, moscas de la fruta)**
Se mantiene	**En grupo**

PROTECCIÓN Y TRÁMITES

CITES	**Apéndice II**
UICN	**En peligro (EN)**

Cuidados de un vistazo

EL TERRARIO	Little is known about keeping *Mantella cowanii*. The species is rare in captivity and is mostly kept without breeding.
TEMPERATURA	The temperature should be 18-21 °C during the day and 16-18 °C at night. This is cooler than for other species in the genus, because the species comes from the highlands at 1,300-2,140 metres above sea level.
HUMEDAD Y PULVERIZACIÓN	Humidity should be 60-80 % (borrowed from *M. baroni*), and the substrate damp but not waterlogged. Mist the enclosure daily with decalcified water.
AGUA Y CALIDAD DEL AGUA	Place a shallow water dish in the enclosure, and keep it full. The frogs are poor swimmers, so the dish must be so shallow that they can easily get out. The depth is not documented; a couple of centimetres is plenty.
ILUMINACIÓN Y UVB	The UVB requirement is not documented for the species, but it is diurnal, so a weak UVB light (Ferguson zone 1) is probably an advantage, provided there is shade and plants so the animals can choose.
ALIMENTACIÓN	In the wild the species feeds on ants, millipedes and mites.
COMPORTAMIENTO Y MANIPULACIÓN	*Mantella cowanii* is diurnal and can be seen patrolling and hunting for food for most of the day. The males call with short, single clicks and defend their territory.
ÁREA DE DISTRIBUCIÓN NATURAL	Madagascar.