herpy

RANAS

Chantaburi Bug-eyed Frog

Theloderma stellatum

Theloderma stellatum is a small, tree-dwelling mossy frog from the lowland rainforest of Thailand, Cambodia, Laos and Vietnam. It resembles a darker version of *Theloderma asperum*, grows to about 3-4 cm and lays its eggs above water in water-filled tree hollows. Its care is similar to that of its relatives, but almost all figures are borrowed from them, because the species is rare in captivity.

Familia: Rhacophoridae

DATOS CLAVE

EL ANIMAL

Tamaño adulto	**3-4 cm** longitud hocico-cloaca
Esperanza de vida	**5-10 años**
Activo	**De noche**
Modo de vida	**Arborícola y semiacuático**
Hábitat	**Selva tropical**
Toxinas cutáneas	**Ninguna conocida**
Dificultad	**Intermedio**

CUIDADOS

Terrario mín.	**45 × 45 × 45 cm** L × An × Al para 4 adultos
Temperatura, día	**20-24 °C**
Temperatura, noche	**17-21 °C**
Humedad	**75-95 %**
Agua	**Zona de agua** profundidad 5-10 cm
UVB	**Opcional** Zona Ferguson 1
Alimentación	**Insectos**
Se mantiene	**En pareja o en grupo**

CRÍA

Reproducción	**Ovípara**
Desarrollo	**Renacuajos en el agua**
Lugar de puesta	**En hojas sobre el agua**
Madurez sexual	**8-10 meses**
Huevos por puesta	**4-15**
Eclosión	**10-14 días**
Periodo larvario	**75-180 días** desde la eclosión hasta la metamorfosis
Alimentación de las larvas	**Alimento completo**
Cuidado parental	**Ninguno**

PROTECCIÓN Y TRÁMITES

CITES	**No incluida**
UICN	**Preocupación menor (LC)**

Cuidados de un vistazo

EL TERRARIO	Little documented knowledge exists on the care of *Theloderma stellatum*.
TEMPERATURA	By day the temperature should be 20-24 °C and at night 17-21 °C.
HUMEDAD Y PULVERIZACIÓN	Humidity should be 75-95 % and never below 60 %, otherwise the frogs may dry out. The large water surface keeps the humidity up on its own, so the frogs do not need much misting.
AGUA Y CALIDAD DEL AGUA	The frog spends a lot of time at the water's edge and even sits under water, so the water section is not there for show.
ILUMINACIÓN Y UVB	The frog is nocturnal and has no documented need for UVB. A weak UVB tube for shade species (Ferguson zone 1) can be used and is optional, but the frogs must always be able to retreat into shade.
ALIMENTACIÓN	*T. stellatum* eats insects. Small crickets are the main food, supplemented with waxworms, flies, springtails for the smallest young, small dubia roaches and soldier fly larvae.
COMPORTAMIENTO Y MANIPULACIÓN	*Theloderma stellatum* is nocturnal and sits hidden by day in tubes, among plants or close to the water. The frog does well in a group as long as there is enough space. The males call at night.
ÁREA DE DISTRIBUCIÓN NATURAL	Tailandia, Camboya, Laos y Vietnam.

La guía completa con mapa y actualizaciones: herpy.org/en/frogs/theloderma-stellatum