herpy

SALAMANDRAS Y TRITONES

Marbled Salamander

Ambystoma opacum

Ambystoma opacum is a small, stocky salamander of 9-11 cm from eastern North America, black with crossbands of white or silvery grey. The species has an unusual breeding biology: mating and egg-laying take place on land in autumn, and the female stays with the eggs until rain fills the hollow and the larvae hatch. The adults live hidden and do best kept cool.

Familia: Ambystomatidae Nombre anterior: *Salamandra armigera; Salamandra fasciata* **Toxinas cutáneas leves**

DATOS CLAVE

EL ANIMAL

Tamaño adulto	**9-11 cm** longitud total con la cola
Esperanza de vida	**4-10 años**
Activo	**De noche**
Modo de vida	**Terrestre y excavador**
Hábitat	**Bosque y prado templados**
Toxinas cutáneas	**Leves** Lávate las manos después del contacto y mantén los dedos alejados de los ojos y la boca
Dificultad	**Principiante**

CRÍA

Reproducción	**Ovípara**
Desarrollo	**Larvas en el agua**
Lugar de puesta	**En el suelo**
Madurez sexual	**12-24 meses**
Huevos por temporada	**50-200**
Puestas por año	**1**
Periodo larvario	**60-270 días** desde la eclosión hasta la metamorfosis
Alimentación de las larvas	**Otros larvas y pequeños invertebrados acuáticos (dafnias, larvas de mosquito)**
Cuidado parental	**Vigila los huevos**

CUIDADOS

Terrario mín.	**40 × 25 × 25 cm** L × An × Al para 1 adulto
Temperatura, día	**18-24 °C**
Temperatura, noche	**16-20 °C**
Humedad	**80-95 %**
Agua	**Bebedero** profundidad 1-2 cm
UVB	**No necesario**
Alimentación	**Insectos y lombrices/caracoles**
Se mantiene	**En solitario o en grupo**

PROTECCIÓN Y TRÁMITES

CITES	**No incluida**
Anexo de la UE	**No incluida**
UICN	**Preocupación menor (LC)**

Cuidados de un vistazo

EL TERRARIO	There is little knowledge about the husbandry of *Ambystoma opacum*, and the advice on substrate and water is partly borrowed from the closely related *Ambystoma maculatum* and *Ambystoma tigrinum*, which are kept far more often.
TEMPERATURA	By day the temperature should be 18-24 °C, and at night it can be a couple of degrees cooler, 16-20 °C. The species tolerates cold better than heat, and above 30 °C for long periods it can die.
HUMEDAD Y PULVERIZACIÓN	The substrate must be damp, but never wet or muddy, because the animal absorbs water through its skin. A damp substrate corresponds to 80-95% humidity. Mist the substrate with chlorine-free water when the surface looks dry.
AGUA Y CALIDAD DEL AGUA	Adults are poor swimmers, and the species drowns if it cannot get out of the water. Deep dishes and water sections must therefore be avoided.
ILUMINACIÓN Y UVB	The species is nocturnal and lives hidden, so it does not need special lighting and does not need UVB either. UV light can damage the eyes and skin of amphibians, and larvae must not be exposed to it.
ALIMENTACIÓN	Adults eat crawling and flying insects, worms and small snails: crickets, earthworms and compost worms, and occasionally waxworms. The feeder animals must be smaller than the animal's head.
COMPORTAMIENTO Y MANIPULACIÓN	Adults lie hidden most of the time and come out at night to hunt. The species does not bite humans. Outside the breeding season the animals live alone, and males in particular bite each other on the snout if they meet.
ÁREA DE DISTRIBUCIÓN NATURAL	Estados Unidos.