herpy

SALAMANDRAS Y TRITONES

Anderson's Crocodile Newt

Echinotriton andersoni

Echinotriton andersoni is a compact, dark brown to black salamander, 14-21 cm long, from the central Ryukyu Islands of Japan. It lives mostly on land in damp forest, has knob-like warts along its sides and can push the tips of its ribs out through the skin as a defence. The species is rare and little known in captivity.

Familia: Salamandridae Nombre anterior: *Pleurodeles andersoni; Tylototriton andersoni* **Toxinas cutáneas leves**

DATOS CLAVE

EL ANIMAL

Tamaño adulto	**14-21 cm** longitud total con la cola
Esperanza de vida	**10-20 años**
Activo	**Al crepúsculo y de noche**
Modo de vida	**Terrestre**
Hábitat	**Bosque seco y humedal**
Toxinas cutáneas	**Leves** Lávate las manos después del contacto y mantén los dedos alejados de los ojos y la boca
Dificultad	**Avanzado**

CRÍA

Reproducción	**Ovípara**
Desarrollo	**Larvas en el agua**
Lugar de puesta	**En el suelo**
Huevos por temporada	**3-46**
Eclosión	**20-27 días**
Periodo larvario	**70-100 días** desde la eclosión hasta la metamorfosis
Alimentación de las larvas	**Otros larvas y pequeños invertebrados acuáticos (dafnias, larvas de mosquito)**
Cuidado parental	**Ninguno**

CUIDADOS

Terrario mín.	**60 × 40 × 40 cm** L × An × Al para 3 adultos
Temperatura, día	**18-25 °C**
Temperatura, noche	**15-20 °C**
Humedad	**60-80 %**
Agua	**Zona de agua** profundidad 10-20 cm
UVB	**No necesario**
Alimentación	**Insectos y lombrices/caracoles**
Se mantiene	**En pareja o en grupo**

PROTECCIÓN Y TRÁMITES

CITES	**Apéndice III**
Anexo de la UE	**Anexo C**
UICN	**Vulnerable (VU)**

Cuidados de un vistazo

EL TERRARIO	Little is known about the care of *Echinotriton andersoni*.
TEMPERATURA	The temperature is 18-25 °C by day and may fall to 15-20 °C at night. It should not stay above 25 °C for long. Heat lamps are unnecessary; keep the terrarium cool in summer and out of direct sun. Winter rest is not documented.
HUMEDAD Y PULVERIZACIÓN	Humidity should be 60-80%. The leaf litter and soil are kept damp but not wet, and part of the substrate should be drier so the animals can choose. Provide a little air exchange so the air does not stand still.
AGUA Y CALIDAD DEL AGUA	The water area is a bowl-shaped depression, about 30 cm in diameter and 10-20 cm deep in the middle, with an even slope all the way round, so the animals are never at risk of drowning.
ILUMINACIÓN Y UVB	The species lives hidden and is most active at dusk and at night. UVB is not necessary. A light period of about 10 hours is suitable, and the light must also be sufficient to keep the plants alive. Avoid direct sun.
ALIMENTACIÓN	The animals take earthworms, crickets, snails, woodlice and small beetles. Earthworms and crickets are suitable as the staple, and the feeder insects must be small enough for the animal to swallow whole.
COMPORTAMIENTO Y MANIPULACIÓN	The animal lives hidden and is most visible at dusk and at night. Groups need plenty of hiding places. The animal is not suited to handling.
ÁREA DE DISTRIBUCIÓN NATURAL	Japón.