herpy

SALAMANDRAS Y TRITONES

California Newt

Taricha torosa

Taricha torosa is a sturdy newt, 13-20 cm long, from the California coast, with a granular brown back and a yellow to orange belly. The skin contains tetrodotoxin, a nerve toxin with no antidote, and the animal can be deadly if swallowed. The species lives mostly on land and enters the water only to breed, and it must be kept cool.

Familia: Salamandridae Nombre anterior: *Cynops torosus; Notophthalmus torosus* **Toxinas cutáneas fuertes**

DATOS CLAVE

EL ANIMAL

Tamaño adulto	**13-20 cm** longitud total con la cola
Esperanza de vida	**12-22 años**
Activo	**De día y de noche**
Modo de vida	**Semiacuático**
Hábitat	**Humedal y bosque y prado templados**
Toxinas cutáneas	**Fuertes** Evita el contacto con la piel y lávate bien las manos después de trabajar en el terrario
Dificultad	**Intermedio**

CRÍA

Reproducción	**Ovípara**
Desarrollo	**Larvas en el agua**
Lugar de puesta	**En el agua**
Madurez sexual	**36 meses**
Huevos por temporada	**7-30**
Eclosión	**14-21 días**
Periodo larvario	**60-90 días** desde la eclosión hasta la metamorfosis
Alimentación de las larvas	**Pequeños invertebrados acuáticos (dafnias, larvas de mosquito)**
Cuidado parental	**Ninguno**

CUIDADOS

Terrario mín.	**60 × 30 × 30 cm** L × An × Al para 2 adultos
Temperatura, día	**16-21 °C**
Temperatura, noche	**14-18 °C**
Humedad	**70-90 %**
Agua	**Bebedero**
UVB	**No necesario**
Alimentación	**Insectos y lombrices/caracoles**
Se mantiene	**En pareja o en grupo**

PROTECCIÓN Y TRÁMITES

CITES	**No incluida**
Anexo de la UE	**No incluida**
UICN	**Casi amenazada (NT)**

Cuidados de un vistazo

EL TERRARIO	There is relatively little knowledge about the care of *Taricha torosa*, so the winter temperatures are borrowed from the closely related *Taricha granulosa*, which is kept more often.
TEMPERATURA	*Taricha torosa* likes the cool. By day the temperature should be 16-21 °C and at night 14-18 °C. The temperature must not exceed 24 °C, and longer periods above 21 °C must be avoided.
HUMEDAD Y PULVERIZACIÓN	The substrate should be damp, but the enclosure should not drip. Humidity is about 70-90%, and daily misting with dechlorinated water keeps the substrate moist.
AGUA Y CALIDAD DEL AGUA	A shallow dish of clean water must always be available, so that the animal can cover itself completely with water. Use tap water treated for chlorine and chloramine; distilled and demineralised water are not used.
ILUMINACIÓN Y UVB	The species does not need UVB. Daylight in the room or a weak fluorescent tube during the day and darkness at night is enough.
ALIMENTACIÓN	*Taricha torosa* is an opportunistic predator that takes almost anything it can overpower. Earthworms are a complete diet in themselves, and compost worms (*Eisenia fetida*) are also taken readily.
COMPORTAMIENTO Y MANIPULACIÓN	*Taricha torosa* is calm, sociable and not aggressive towards others of its kind of the same size.
ÁREA DE DISTRIBUCIÓN NATURAL	Estados Unidos.

La guía completa con mapa y actualizaciones: herpy.org/en/salamanders/taricha-torosa