herpy

SALAMANDRAS Y TRITONES

Danube Crested Newt

Triturus dobrogicus

Triturus dobrogicus is a slender crested newt, 13-18 cm long, from the Danube basin and the plains of the large rivers. It is the smallest and most aquatic of the crested newts, with an orange to red belly and sharply defined black spots. The male develops a serrated crest in spring. The species is rare in captivity, and its care follows that of the other crested newts.

Familia: Salamandridae **Toxinas cutáneas leves**

DATOS CLAVE

EL ANIMAL

Tamaño adulto	**13-18 cm** longitud total con la cola
Esperanza de vida	**7-17 años**
Activo	**Al crepúsculo y de noche**
Modo de vida	**Semiacuático**
Hábitat	**Humedal y bosque y prado templados**
Toxinas cutáneas	**Leves** Lávate las manos después del contacto y mantén los dedos alejados de los ojos y la boca
Dificultad	**Intermedio**

CUIDADOS

Terrario mín.	**45 × 30 × 30 cm** L × An × Al para 4 adultos
Temperatura, día	**12-20 °C**
Temperatura, noche	**8-15 °C**
Humedad	**70-95 %**
Agua	**Acuaterrario** profundidad 15-30 cm
Temperatura del agua	**10-20 °C**
UVB	**No necesario**
Reposo invernal	**4-10 °C** 6-12 semanas
Alimentación	**Insectos y lombrices/caracoles**
Se mantiene	**En pareja o en grupo**

CRÍA

Reproducción	**Ovípara**
Desarrollo	**Larvas en el agua**
Lugar de puesta	**En el agua**
Madurez sexual	**12-36 meses**
Huevos por temporada	**200**
Eclosión	**14-35 días**
Periodo larvario	**60-120 días** desde la eclosión hasta la metamorfosis
Alimentación de las larvas	**Alimento completo, otros larvas y pequeños invertebrados acuáticos (dafnias, larvas de mosquito)**
Cuidado parental	**Ninguno**

PROTECCIÓN Y TRÁMITES

CITES	**No incluida**
Anexo de la UE	**No incluida**
UICN	**Preocupación menor (LC)**

Cuidados de un vistazo

EL TERRARIO	Little is known about the care of *Triturus dobrogicus*, and the species is rare in captivity.
TEMPERATURA	*Triturus dobrogicus* loves cool conditions. The water is kept at 10-20 °C and must never become warmer; the animals are active down to about 5 °C.
HUMEDAD Y PULVERIZACIÓN	In the aquatic phase humidity is not a problem. In the land phase the soft skin dries out quickly, so the substrate must be damp, with a humidity of 70-95%.
AGUA Y CALIDAD DEL AGUA	Water depth is 15-30 cm, and the water is kept at 10-20 °C. The species is used to river water and tolerates many kinds of water, but chlorine harms it; tap water that has stood for a couple of days is unproblematic.
ILUMINACIÓN Y UVB	The animals are nocturnal and do not need UVB. The plants need light to grow, and an ordinary aquarium lamp is enough.
ALIMENTACIÓN	In the water *T. dobrogicus* takes earthworms, freshwater shrimp (*Gammarus*), bloodworms, daphnia and white worms. Frozen bloodworms and fish food are also taken, but often have to be offered with tweezers at first.
COMPORTAMIENTO Y MANIPULACIÓN	*Triturus dobrogicus* is nocturnal and hides in vegetation by day. In the water the animals snap at anything that moves, including each other during feeding, and in the wild they also eat smaller newts and tadpoles.
ÁREA DE DISTRIBUCIÓN NATURAL	Austria, Chequia, Eslovaquia, Hungría, Eslovenia, Croacia, Bosnia y Herzegovina, Serbia, Rumanía, Ucrania, Bulgaria y Moldavia.