herpy

SALAMANDRAS Y TRITONES

Southern Crested Newt

Triturus karelinii

Triturus karelinii is a large, robust crested newt of 12-18 cm from the Crimea, the Caucasus and the southern shore of the Caspian Sea. It has brown to grey, warty skin, and the throat and belly are orange or yellow with small black spots. In spring the male develops a tall, serrated crest. The species is hardy and easy to keep, and it can be kept both on land and in water.

Familia: Salamandridae Nombre anterior: *Triturus cristatus karelinii* **Toxinas cutáneas leves**

DATOS CLAVE

EL ANIMAL

Tamaño adulto	**12-18 cm** longitud total con la cola
Esperanza de vida	**8-11 años**
Activo	**Al crepúsculo y de noche**
Modo de vida	**Semiacuático**
Hábitat	**Humedal y bosque y prado templados**
Toxinas cutáneas	**Leves** Lávate las manos después del contacto y mantén los dedos alejados de los ojos y la boca
Dificultad	**Principiante**

CUIDADOS

Terrario mín.	**45 × 30 × 30 cm** L × An × Al para 4 adultos
Temperatura, día	**12-20 °C**
Temperatura, noche	**8-15 °C**
Humedad	**70-95 %**
Agua	**Acuaterrario** profundidad 15-30 cm
Temperatura del agua	**10-20 °C**
UVB	**No necesario**
Reposo invernal	**4-10 °C** 6-12 semanas
Alimentación	**Insectos y lombrices/caracoles**
Se mantiene	**En pareja o en grupo**

CRÍA

Reproducción	**Ovípara**
Desarrollo	**Larvas en el agua**
Lugar de puesta	**En el agua**
Madurez sexual	**12-36 meses**
Huevos por temporada	**150-400**
Eclosión	**14-35 días**
Periodo larvario	**56-112 días** desde la eclosión hasta la metamorfosis
Alimentación de las larvas	**Alimento completo, otros larvas y pequeños invertebrados acuáticos (dafnias, larvas de mosquito)**
Cuidado parental	**Ninguno**

PROTECCIÓN Y TRÁMITES

CITES	**No incluida**
Anexo de la UE	**No incluida**
UICN	**Preocupación menor (LC)**

Cuidados de un vistazo

EL TERRARIO	*Triturus karelinii* can be kept following its natural cycle, with an aquatic phase and a terrestrial phase, or in water all year.
TEMPERATURA	The water should be kept below 25 °C, because in warmer water the animals try to leave it. The animals are fully active down to about 10 °C.
HUMEDAD Y PULVERIZACIÓN	In the aquatic phase humidity is not an issue. In the terrestrial phase the soft skin dries out quickly, so the substrate must be kept damp, with a humidity of 70-95%.
AGUA Y CALIDAD DEL AGUA	A water depth of 15 cm is enough, and deeper is fine: 15-30 cm gives the animals room to swim. The water is kept at 10-20 °C.
ILUMINACIÓN Y UVB	The animals are nocturnal and do not need UVB or artificial light for their own sake. The plants need light to grow, and an ordinary aquarium lamp is enough.
ALIMENTACIÓN	In the aquatic phase *T. karelinii* takes a wide range of food: earthworms, freshwater shrimp (*Gammarus*), bloodworms, daphnia and white worms.
COMPORTAMIENTO Y MANIPULACIÓN	*Triturus karelinii* is robust and easy to keep. In the water the animals snap hard at anything that moves, including each other during feeding, and in the wild crested newts also eat smaller amphibians.
ÁREA DE DISTRIBUCIÓN NATURAL	Ucrania, Rusia, Georgia, Azerbaiyán y Irán.