herpy

SERPIENTES

Brown House Snake

Boaedon capensis

Boaedon capensis is a slender, nocturnal constrictor from southern Africa, where it lives in scrub, grassland and around human dwellings and hunts rodents and lizards. In the enclosure it is hardy and simple to keep, and adults usually become calm in the hand, but the species is a skilled escape artist, and juveniles are fast.

Familia: Lamprophiidae Nombre anterior: *Lamprophis capensis; Boaedon capense*

DATOS CLAVE

EL ANIMAL

Tamaño adulto	**60-120 cm** longitud total con la cola
Esperanza de vida	**10-20 años**
Activo	**De noche**
Modo de vida	**Terrestre**
Hábitat	**Semidesierto y sabana**
Dificultad	**Principiante**

CUIDADOS

Terrario mín.	**90 × 45 × 40 cm** L × An × Al para 1 adulto
Temperatura, día	**24-28 °C**
Punto caliente	**30-32 °C**
Temperatura, noche	**20-24 °C**
Humedad	**40-60 %**
Agua	**Bebedero**
UVB	**Opcional**
Alimentación	**Vertebrados**
Se mantiene	**En solitario**

CRÍA

Reproducción	**Ovípara**
Tamaño de la puesta	**5-16 huevos**
Puestas por año	**1-4**
Incubación	**55-65 días** a 29-30 °C

PROTECCIÓN Y TRÁMITES

CITES	**No incluida**
Anexo de la UE	**No incluida**
UICN	**Preocupación menor (LC)**

Cuidados de un vistazo

EL TERRARIO

An adult *B. capensis* needs an enclosure of at least 90 × 45 × 40 cm. Larger is always better, and large females do well in 100 × 50 × 40 cm.

TEMPERATURA

The enclosure must have a clear temperature gradient. During the day it is 24-28 °C, with about 24 °C at the cool end. The warmest spot, where the snake can lie and warm itself, should be 30-32 °C.

HUMEDAD Y AGUA

Humidity should be 40-60%. In most rooms this is achieved with a water bowl, and it is rarely necessary to mist.

ILUMINACIÓN Y UVB

B. capensis is nocturnal and does not need bright light. A fixed light rhythm of about 12 hours of light and 12 hours of dark all year round gives a clear daily rhythm. UVB is not necessary for animals that are fed whole prey.

ALIMENTACIÓN

Boaedon capensis kills its prey by constriction, has a strong feeding response and usually eats readily. It is fed whole, thawed mice and, for large females, small rats, never live prey.

COMPORTAMIENTO Y MANIPULACIÓN

Juveniles and subadults can be fast, nippy and restless, and they can spring out of the enclosure like a released spring when the lid is opened. Open the lid carefully over a safe spot.

ÁREA DE DISTRIBUCIÓN NATURAL

Sudáfrica, Lesoto, Esuatini, Botsuana, Zimbabue, Mozambique, Zambia y Malaui.