herpy

SERPIENTES

Plains Garter Snake

Thamnophis radix

Thamnophis radix is a slim, striped garter snake from the prairies and open grasslands of central North America. It has a yellow or orange stripe down the back and grows to 40-70 cm. It is very tolerant of cold and gives birth to live young. It is hardy and can be kept on thawed mice, but it escapes through the smallest gaps.

Familia: Colubridae

DATOS CLAVE

EL ANIMAL

Tamaño adulto	**40-70 cm** longitud total con la cola
Esperanza de vida	**5-10 años**
Activo	**De día**
Modo de vida	**Terrestre**
Hábitat	**Humedal y bosque y prado templados**
Dificultad	**Principiante**

CUIDADOS

Terrario mín.	**100 × 50 × 50 cm** L × An × Al para 1 adulto
Temperatura, día	**23-27 °C**
Punto caliente	**29-32 °C**
Temperatura, noche	**18-24 °C**
Humedad	**35-60 %**
Agua	**Bebedero**
UVB	**Recomendado** Zona Ferguson 2
Reposo invernal	**8-15 °C** 8-16 semanas
Alimentación	**Vertebrados y lombrices/caracoles**
Se mantiene	**En solitario o en grupo**

CRÍA

Reproducción	**Vivípara**
Madurez sexual	**24-36 meses**
Tamaño de la camada	**5-60 crías**
Puestas por año	**1**
Gestación	**83-102 días**

PROTECCIÓN Y TRÁMITES

CITES	**No incluida**
Anexo de la UE	**No incluida**
UICN	**Preocupación menor (LC)**

Cuidados de un vistazo

EL TERRARIO	Little species-specific knowledge exists about keeping *Thamnophis radix*.
TEMPERATURA	There must be a clear temperature gradient. The basking spot should be 29-32 °C measured on the surface, the cool end 23-27 °C and the night 18-24 °C.
HUMEDAD Y AGUA	The air humidity should be 35-60 %, highest at the cool end and at night. The enclosure should be dry, because persistently damp conditions cause blisters on the belly and fungal infections.
ILUMINACIÓN Y UVB	*Thamnophis radix* is diurnal and thrives in strong daylight. UVB is not strictly necessary, but is recommended.
ALIMENTACIÓN	In the wild *Thamnophis radix* eats earthworms, slugs, frogs, toads, salamanders, fish and the occasional mouse.
COMPORTAMIENTO Y MANIPULACIÓN	*Thamnophis radix* is quick and alert, but usually becomes calm if handled regularly from a young age. Let new animals settle in for about two weeks, and start handling them only when they are feeding steadily.
ÁREA DE DISTRIBUCIÓN NATURAL	Canadá y Estados Unidos.