herpy

SALAMANDRES ET TRITONS

Southern Crested Newt

Triturus karelinii

Triturus karelinii is a large, robust crested newt of 12-18 cm from the Crimea, the Caucasus and the southern shore of the Caspian Sea. It has brown to grey, warty skin, and the throat and belly are orange or yellow with small black spots. In spring the male develops a tall, serrated crest. The species is hardy and easy to keep, and it can be kept both on land and in water.

Famille: Salamandridae Ancien nom: *Triturus cristatus karelinii* **Toxines cutanées faibles**

INFORMATIONS ESSENTIELLES

L'ANIMAL

Taille adulte	**12-18 cm** longueur totale queue comprise
Espérance de vie	**8-11 ans**
Activité	**Au crépuscule et la nuit**
Mode de vie	**Semi-aquatique**
Habitat	**Zone humide et forêt et prairie tempérées**
Toxines cutanées	**Faibles** Lavez-vous les mains après le contact et gardez les doigts loin des yeux et de la bouche
Difficulté	**Débutant**

ENTRETIEN

Terrarium min.	**45 × 30 × 30 cm** L × l × H pour 4 adultes
Température, jour	**12-20 °C**
Température, nuit	**8-15 °C**
Hygrométrie	**70-95 %**
Eau	**Aquaterrarium** profondeur 15-30 cm
Température de l'eau	**10-20 °C**
UVB	**Non nécessaires**
Repos hivernal	**4-10 °C** 6-12 semaines
Alimentation	**Insectes et vers/escargots**
Détention	**En couple ou en groupe**

REPRODUCTION

Reproduction	**Ovipare**
Développement	**Larves dans l'eau**
Lieu de ponte	**Dans l'eau**
Maturité sexuelle	**12-36 mois**
Œufs par saison	**150-400**
Éclosion	**14-35 jours**
Durée du stade larvaire	**56-112 jours** de l'éclosion à la métamorphose
Alimentation des larves	**Aliment complet, autres larves et petits invertébrés aquatiques (daphnies, larves de moustiques)**
Soins parentaux	**Aucun**

PROTECTION ET FORMALITÉS

CITES	**Non inscrite**
Annexe UE	**Non inscrite**
UICN	**Préoccupation mineure (LC)**

L'entretien en bref

LE TERRARIUM	*Triturus karelinii* can be kept following its natural cycle, with an aquatic phase and a terrestrial phase, or in water all year.
TEMPÉRATURE	The water should be kept below 25 °C, because in warmer water the animals try to leave it. The animals are fully active down to about 10 °C.
HYGROMÉTRIE ET BRUMISATION	In the aquatic phase humidity is not an issue. In the terrestrial phase the soft skin dries out quickly, so the substrate must be kept damp, with a humidity of 70-95%.
EAU ET QUALITÉ DE L'EAU	A water depth of 15 cm is enough, and deeper is fine: 15-30 cm gives the animals room to swim. The water is kept at 10-20 °C.
ÉCLAIRAGE ET UVB	The animals are nocturnal and do not need UVB or artificial light for their own sake. The plants need light to grow, and an ordinary aquarium lamp is enough.
ALIMENTATION	In the aquatic phase *T. karelinii* takes a wide range of food: earthworms, freshwater shrimp (*Gammarus*), bloodworms, daphnia and white worms.
COMPORTEMENT ET MANIPULATION	*Triturus karelinii* is robust and easy to keep. In the water the animals snap hard at anything that moves, including each other during feeding, and in the wild crested newts also eat smaller amphibians.
AIRE D'ORIGINE	Ukraine, Russie, Géorgie, Azerbaïdjan et Iran.