herpy

RANE FRECCIA

Niceforo's Poison Frog

Ameerega ingeri

Ameerega ingeri is a small, diurnal poison dart frog of 20-27 mm from the foothills of the Andes in southern Colombia. The back is uniformly dark, the flanks have dark blue spots, and there are bright orange spots in the armpits and on the calves. The species is hardly ever kept, so care is based on the climate of its range and on related species.

Famiglia: Dendrobatidae Nome precedente: *Dendrobates ingeri; Epipedobates ingeri; Phyllobates ingeri*

DATI ESSENZIALI

L'ANIMALE

Dimensioni da adulto	**20-27 mm** lunghezza muso-cloaca
Aspettativa di vita	**10-15 anni**
Attività	**Di giorno**
Stile di vita	**Terrestre**
Habitat	**Foresta pluviale**
Tossine cutanee in natura	**Lievi**
Tossine cutanee in cattività	**Nessuna** Animali nati in cattività. Gli animali catturati in natura possono conservare le tossine a lungo
Difficoltà	**Intermedio**

CURA

Terrario min.	**60 × 40 × 40 cm** L × P × H per 2 adulti
Temperatura, giorno	**24-26 °C**
Temperatura, notte	**20-22 °C**
Umidità	**70-100%**
Acqua	**Ciotola d'acqua** profondità 1-2 cm
UVB	**Facoltativi** Zona Ferguson 1
Alimentazione	**Piccoli invertebrati (collemboli, moscerini della frutta)**
Detenzione	**In coppia**

RIPRODUZIONE

Riproduzione	**Ovipara**
Sviluppo	**Girini trasportati fino all'acqua**
Luogo di deposizione	**Sul terreno**
Uova per deposizione	**4-16**
Schiusa	**10-22 giorni**
Periodo larvale	**70-90 giorni** dalla schiusa alla metamorfosi
Alimentazione delle larve	**Alghe/piante e mangime completo**
Cure parentali	**Trasporta i girini**

TUTELA E DOCUMENTI

CITES	**Appendice II**
IUCN	**Dati insufficienti (DD)**

La cura in breve

IL TERRARIO	Almost nothing is known about the care of *Ameerega ingeri*, and the species is hardly ever kept.
TEMPERATURA	Keep the enclosure at 24-26 °C by day, and let the temperature fall to 20-22 °C at night. The temperature must not exceed 28 °C, because frogs tolerate heat poorly.
UMIDITÀ E NEBULIZZAZIONE	Humidity should be 70-80 % in the middle of the day and close to 100 % in the morning and evening. A rain system or a fogger with a timer is easiest; otherwise mist two to three times a day with reverse-osmosis water or rainwater.
ACQUA E QUALITÀ DELL'ACQUA	Provide a shallow water dish with 1-2 cm of water, so that the frogs can easily climb out of it, and with stones or moss at the edge. Use reverse-osmosis water, rainwater or dechlorinated tap water, and change it often.
ILLUMINAZIONE E UVB	The lighting is primarily for the plants: 10-12 hours a day with LED lighting with a plant-suitable spectrum. It is not documented that *Ameerega ingeri* needs UVB.
ALIMENTAZIONE	In the wild *Ameerega ingeri* lives mostly on small ants and other small arthropods from the leaf litter.
COMPORTAMENTO E MANIPOLAZIONE	*Ameerega ingeri* is diurnal and most active in the morning. It lives on the ground in the leaf litter, but likes to sleep on leaves some way above the ground, so plants with broad leaves are an advantage.
AREALE D'ORIGINE	Colombia.