herpy

SALAMANDRE E TRITONI

California Newt

Taricha torosa

Taricha torosa is a sturdy newt, 13-20 cm long, from the California coast, with a granular brown back and a yellow to orange belly. The skin contains tetrodotoxin, a nerve toxin with no antidote, and the animal can be deadly if swallowed. The species lives mostly on land and enters the water only to breed, and it must be kept cool.

Famiglia: Salamandridae Nome precedente: *Cynops torosus; Notophthalmus torosus* **Tossine cutanee forti**

DATI ESSENZIALI

L'ANIMALE

Dimensioni da adulto	**13-20 cm** lunghezza totale inclusa la coda
Aspettativa di vita	**12-22 anni**
Attività	**Di giorno e di notte**
Stile di vita	**Semiacquatico**
Habitat	**Zona umida e bosco e prato temperati**
Tossine cutanee	**Forti** Evita il contatto con la pelle e lavati accuratamente le mani dopo aver lavorato nel terrario
Difficoltà	**Intermedio**

CURA

Terrario min.	**60 × 30 × 30 cm** L × P × H per 2 adulti
Temperatura, giorno	**16-21 °C**
Temperatura, notte	**14-18 °C**
Umidità	**70-90%**
Acqua	**Ciotola d'acqua**
UVB	**Non necessari**
Alimentazione	**Insetti e vermi/lumache**
Detenzione	**In coppia o in gruppo**

RIPRODUZIONE

Riproduzione	**Ovipara**
Sviluppo	**Larve in acqua**
Luogo di deposizione	**In acqua**
Maturità sessuale	**36 mesi**
Uova per stagione	**7-30**
Schiusa	**14-21 giorni**
Periodo larvale	**60-90 giorni** dalla schiusa alla metamorfosi
Alimentazione delle larve	**Piccoli invertebrati acquatici (dafnie, larve di zanzara)**
Cure parentali	**Nessuna**

TUTELA E DOCUMENTI

CITES	**Non elencata**
Allegato UE	**Non elencata**
IUCN	**Quasi minacciata (NT)**

La cura in breve

IL TERRARIO	There is relatively little knowledge about the care of *Taricha torosa*, so the winter temperatures are borrowed from the closely related *Taricha granulosa*, which is kept more often.
TEMPERATURA	*Taricha torosa* likes the cool. By day the temperature should be 16-21 °C and at night 14-18 °C. The temperature must not exceed 24 °C, and longer periods above 21 °C must be avoided.
UMIDITÀ E NEBULIZZAZIONE	The substrate should be damp, but the enclosure should not drip. Humidity is about 70-90%, and daily misting with dechlorinated water keeps the substrate moist.
ACQUA E QUALITÀ DELL'ACQUA	A shallow dish of clean water must always be available, so that the animal can cover itself completely with water. Use tap water treated for chlorine and chloramine; distilled and demineralised water are not used.
ILLUMINAZIONE E UVB	The species does not need UVB. Daylight in the room or a weak fluorescent tube during the day and darkness at night is enough.
ALIMENTAZIONE	*Taricha torosa* is an opportunistic predator that takes almost anything it can overpower. Earthworms are a complete diet in themselves, and compost worms (*Eisenia fetida*) are also taken readily.
COMPORTAMENTO E MANIPOLAZIONE	*Taricha torosa* is calm, sociable and not aggressive towards others of its kind of the same size.
AREALE D'ORIGINE	Stati Uniti.

La guida completa con mappa e aggiornamenti: herpy.org/en/salamanders/taricha-torosa