herpy

SALAMANDRE E TRITONI

Southern Crested Newt

Triturus karelinii

Triturus karelinii is a large, robust crested newt of 12-18 cm from the Crimea, the Caucasus and the southern shore of the Caspian Sea. It has brown to grey, warty skin, and the throat and belly are orange or yellow with small black spots. In spring the male develops a tall, serrated crest. The species is hardy and easy to keep, and it can be kept both on land and in water.

Famiglia: Salamandridae Nome precedente: *Triturus cristatus karelinii* **Tossine cutanee lievi**

DATI ESSENZIALI

L'ANIMALE

Dimensioni da adulto	**12-18 cm** lunghezza totale inclusa la coda
Aspettativa di vita	**8-11 anni**
Attività	**Al crepuscolo e di notte**
Stile di vita	**Semiacquatico**
Habitat	**Zona umida e bosco e prato temperati**
Tossine cutanee	**Lievi** Lavati le mani dopo il contatto e tieni le dita lontane da occhi e bocca
Difficoltà	**Principiante**

CURA

Terrario min.	**45 × 30 × 30 cm** L × P × H per 4 adulti
Temperatura, giorno	**12-20 °C**
Temperatura, notte	**8-15 °C**
Umidità	**70-95%**
Acqua	**Acquaterrario** profondità 15-30 cm
Temperatura dell'acqua	**10-20 °C**
UVB	**Non necessari**
Riposo invernale	**4-10 °C** 6-12 settimane
Alimentazione	**Insetti e vermi/lumache**
Detenzione	**In coppia o in gruppo**

RIPRODUZIONE

Riproduzione	**Ovipara**
Sviluppo	**Larve in acqua**
Luogo di deposizione	**In acqua**
Maturità sessuale	**12-36 mesi**
Uova per stagione	**150-400**
Schiusa	**14-35 giorni**
Periodo larvale	**56-112 giorni** dalla schiusa alla metamorfosi
Alimentazione delle larve	**Mangime completo, altri larve e piccoli invertebrati acquatici (dafnie, larve di zanzara)**
Cure parentali	**Nessuna**

TUTELA E DOCUMENTI

CITES	**Non elencata**
Allegato UE	**Non elencata**
IUCN	**Minima preoccupazione (LC)**

La cura in breve

IL TERRARIO	*Triturus karelinii* can be kept following its natural cycle, with an aquatic phase and a terrestrial phase, or in water all year.
TEMPERATURA	The water should be kept below 25 °C, because in warmer water the animals try to leave it. The animals are fully active down to about 10 °C.
UMIDITÀ E NEBULIZZAZIONE	In the aquatic phase humidity is not an issue. In the terrestrial phase the soft skin dries out quickly, so the substrate must be kept damp, with a humidity of 70-95%.
ACQUA E QUALITÀ DELL'ACQUA	A water depth of 15 cm is enough, and deeper is fine: 15-30 cm gives the animals room to swim. The water is kept at 10-20 °C.
ILLUMINAZIONE E UVB	The animals are nocturnal and do not need UVB or artificial light for their own sake. The plants need light to grow, and an ordinary aquarium lamp is enough.
ALIMENTAZIONE	In the aquatic phase *T. karelinii* takes a wide range of food: earthworms, freshwater shrimp (*Gammarus*), bloodworms, daphnia and white worms.
COMPORTAMENTO E MANIPOLAZIONE	*Triturus karelinii* is robust and easy to keep. In the water the animals snap hard at anything that moves, including each other during feeding, and in the wild crested newts also eat smaller amphibians.
AREALE D'ORIGINE	Ucraina, Russia, Georgia, Azerbaigian e Iran.