herpy

AGAMEN

Green Crested Lizard

Bronchocela cristatella

Bronchocela cristatella is a pale green, very slender arboreal agamid from the forests of South-East Asia, where the tail makes up over three quarters of the length. It reaches 50-58 cm, is diurnal and hunts insects in the trees. It can change from clear green to brown when it becomes agitated, and it is fast, a strong jumper and shy.

Familie: Agamidae Vroegere naam: *Calotes cristatellus; Agama cristatella; Agama moluccana*

KERNGEGEVENS

HET DIER

Volwassen grootte	**50-58 cm** totale lengte incl. staart
Levensduur	**5-10 jaar**
Actief	**Overdag**
Levenswijze	**Boombewonend**
Leefgebied	**Regenwoud**
Moeilijkheidsgraad	**Ervaren**

VERZORGING

Min. terrarium	**120 × 60 × 150 cm** L × B × H voor 1 volwassen dier
Temperatuur overdag	**25-30 °C**
Zonneplek	**32-38 °C**
Temperatuur 's nachts	**22-25 °C**
Luchtvochtigheid	**70-90%**
Water	**Geen waterbak**
UVB	**Aanbevolen** Fergusonzone 2
Voeding	**Insecten**
Gehouden	**Solitair of in paren**

KWEEK

Voortplanting	**Eierleggend**
Legselgrootte	**2 eieren**
Incubatie	**47-80 dagen** bij 27-31 °C

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
EU-bijlage	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	Little is known about the care of *Bronchocela cristatella*.
TEMPERATUUR	The temperature follows the climate in the range. The daytime temperature in the enclosure should be 25-30 °C, and at a basking spot high up on a branch there should be 32-38 °C.
LUCHTVOCHTIGHEID EN WATER	Relative humidity should be 70-90 %, as in a tropical lowland forest. Mist the enclosure thoroughly twice a day so that the leaves and branches are wet. The animal drinks droplets from the plants, and a bowl is not necessary.
VERLICHTING EN UVB	The species is diurnal and basks at forest edges and in clearings, so it needs strong light and UVB.
VOEDING	*Bronchocela cristatella* feeds on insects, which it catches in bushes and trees, including ticks.
GEDRAG EN OMGANG	The species is shy and easily stressed. Under stress it turns brown, and it may flee into the foliage or jump from the branches. It is unsuitable for handling and is an animal to look at.
NATUURLIJK VERSPREIDINGSGEBIED	Maleisië, Singapore, Indonesië, Filipijnen, Thailand en Brunei.