herpy

AGAMEN

Sudan Mastigure

Uromastyx dispar

Uromastyx dispar is a spiny-tailed lizard from the Sahara and Sahel with four subspecies, including the large *U. d. flavifasciata* and the frequently kept *U. d. maliensis* from Mali. The species grows to 28-50 cm depending on form, is herbivorous and digs down into a deep substrate. It needs a large, hot and dry enclosure with powerful UVB and is more an animal to watch than to handle.

Familie: Agamidae Vroegere naam: *Uromastyx flavifasciata; Uromastyx maliensis*

KERNGEGEVENS

HET DIER

Volwassen grootte	**28-50 cm** totale lengte incl. staart
Gewicht	**200-400 g**
Levensduur	**15-25 jaar**
Actief	**Overdag**
Levenswijze	**Rotsbewonend, bodembewonend en gravend**
Leefgebied	**Woestijn en rotsachtig terrein**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**120 × 60 × 60 cm** L × B × H voor 1 volwassen dier
Temperatuur overdag	**29-35 °C**
Zonneplek	**45-50 °C**
Temperatuur ’s nachts	**20-22 °C**
Luchtvochtigheid	**20-30%**
Water	**Geen waterbak**
UVB	**Onmisbaar** Fergusonzone 3
Winterrust	**18-21 °C** 8-12 weken
Voeding	**Planten**
Gehouden	**Solitair of in paren**

KWEEK

Voortplanting	**Eierleggend**
Geslachtsrijp	**36 maanden**
Legselgrootte	**10-20 eieren**
Legsels per jaar	**1**
Incubatie	**60-80 dagen** bij 33 °C

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
EU-bijlage	**Bijlage B**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM

An adult of the nominate form or *U. d. maliensis* needs an enclosure of at least 120 × 60 × 60 cm (length × width × height), and bigger is better. The larger *U. d. flavifasciata* needs at least 150 × 60 × 60 cm.

TEMPERATUUR

The basking spot must have a surface temperature of 45-50 °C, measured with an infrared thermometer directly under the lamp. The middle of the enclosure is at 35-38 °C and the cool end at 29-35 °C.

LUCHTVOCHTIGHEID EN WATER

Humidity must be below 30 %. The species tolerates slightly higher humidity as long as the enclosure is dry and warm enough, but it does not tolerate being cold and damp. Do not mist the enclosure.

VERLICHTING EN UVB

The species is diurnal and needs strong light. Use an LED lamp or a fluorescent tube of about 6500 K as daylight in addition to the heat lamp.

VOEDING

Uromastyx dispar is herbivorous.

GEDRAG EN OMGANG

Adults are territorial, especially males towards males, and two males must never be kept together.

NATUURLIJK VERSPREIDINGSGEBIED

Algerije, Westelijke Sahara, Mali, Mauritanië, Soedan en Tsjaad.