herpy

GEKKO'S

Wiegmann's striped gecko

Gonatodes vittatus

A small, diurnal gecko from Venezuela, Colombia and the islands off the coast, where the male has a white stripe with a black edge down the back and a yellow head. It climbs on tree trunks and walls but cannot climb glass, and it does well in a small enclosure if kept alone or as a pair.

Familie: Sphaerodactylidae Vroegere naam: *Gymnodactylus vittatus; Gonatodes gillii*

KERNGEGEVENS

HET DIER

Volwassen grootte	**7-8 cm** totale lengte incl. staart
Levensduur	**10-20 jaar**
Actief	**Overdag**
Levenswijze	**Boombewonend en bodembewonend**
Leefgebied	**Droog bos**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**30 × 30 × 30 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**24-27 °C**
Zonneplek	**29-32 °C**
Temperatuur 's nachts	**20-24 °C**
Luchtvochtigheid	**50-60%**
UVB	**Optioneel**
Voeding	**Insecten en kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**Solitair of in paren**

KWEEK

Voortplanting	**Eierleggend**
Legselgrootte	**1 eieren**
Incubatie	**65-110 dagen** bij 25-30 °C

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
EU-bijlage	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM
A pair needs an enclosure of at least 30 × 30 × 30 cm (length × width × height), and larger is better, because more space gives more calm.

TEMPERATUUR
By day the enclosure should be 24-27 °C. A warm spot of 29-32 °C can be made with a heat mat or a lamp, but it is not necessary if the temperature is otherwise high and stable.

LUCHTVOCHTIGHEID EN WATER
Gonatodes vittatus thrives in drier air than most tropical geckos. Keep the humidity at 50-60 %.

VERLICHTING EN UVB
The species is diurnal but does not bask, and UVB is optional. Weak UV light can make it easier to get the animals to breed, but too much UV can cause problems with shedding.

VOEDING
Gonatodes vittatus eats insects and other small arthropods.

GEDRAG EN OMGANG
The species is bold and curious and is often seen during the day, but it is not an animal to be handled. The skin can tear if the animal is grabbed, and the animals are easily stressed.

NATUURLIJK VERSPREIDINGSGEBIED
Colombia, Venezuela, Guyana, Trinidad en Tobago, Aruba en Curaçao.