herpy

GEKKO'S

Banded velvet gecko

Homopholis fasciata

A sturdy, velvety gecko from the savannas of East Africa with dark bands across the back and both adhesive lamellae and claws on the toes. It is most active at night, hides in holes in large trees and plays dead when threatened. Care is described only in broad outline.

Familie: Gekkonidae Vroegere naam: *Platypholis fasciata; Homopholis erlangeri*

KERNGEGEVENS

HET DIER

Volwassen grootte	**9-11 cm** totale lengte incl. staart
Levensduur	**8-12 jaar**
Actief	**In de schemering en 's nachts**
Levenswijze	**Boombewonend**
Leefgebied	**Savanne**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**40 × 40 × 60 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**24-28 °C**
Zonneplek	**28-32 °C**
Temperatuur 's nachts	**19-24 °C**
Luchtvochtigheid	**60-80%**
UVB	**Optioneel**
Winterrust	**19-23 °C** 12-16 weken
Voeding	**Insecten**
Gehouden	**In paren of één mannetje met meerdere vrouwtjes**

KWEEK

Voortplanting	**Eierleggend**
Legselgrootte	**2 eieren**
Legsels per jaar	**4-5**
Incubatie	**60-80 dagen** bij 26-29 °C

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
EU-bijlage	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM

Homopholis fasciata climbs, so a pair needs a tall enclosure of at least 40 × 40 × 60 cm (length × width × height). Larger is always better, especially if you keep more than two animals.

TEMPERATUUR

By day the temperature should be 24-28 °C, and there should be a warmer spot of 28-32 °C, for example under a small 10-20 W halogen bulb or over a heat mat. At night it may fall 3-6 °C, to about 19-24 °C.

LUCHTVOCHTIGHEID EN WATER

Humidity should be moderate to high, 60-80 %, with a gradient from one end of the enclosure to the other.

VERLICHTING EN UVB

The species is active at dusk and at night and can be seen in daylight, but hides in the shade in the middle of the day. Use ordinary plant lighting on a timer, 12 hours on and 12 hours off.

VOEDING

The staple food is crickets, supplemented with cockroaches, mealworms and other insects of suitable size, so that the diet is varied. Waxworms can be given occasionally.

GEDRAG EN OMGANG

Homopholis fasciata can bite and hold on, and it plays dead and often lies motionless when frightened, so do not pick it up without reason. Give new animals time to settle in.

NATUURLIJK VERSPREIDINGSGEBIED

Ethiopië, Kenia, Somalië en Tanzania.