herpy

GEKKO'S

Cameroon dwarf gecko

Lygodactylus conraui

A small, diurnal climber from the West African forests, and the only green gecko in the region. It sits on tree trunks, palm fronds and house walls at shady forest edges. It is rare in vivaria, and its care is based on closely related dwarf day geckos.

Familie: Gekkonidae

KERNGEGEVENS

HET DIER

Volwassen grootte	**7-9 cm** totale lengte incl. staart
Levensduur	**5-8 jaar**
Actief	**Overdag**
Levenswijze	**Boombewonend**
Leefgebied	**Regenwoud**
Moeilijkheidsgraad	**Ervaren**

VERZORGING

Min. terrarium	**30 × 30 × 45 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**24-27 °C**
Zonneplek	**32-34 °C**
Temperatuur 's nachts	**21-24 °C**
Luchtvochtigheid	**60-80%**
UVB	**Aanbevolen** Fergusonzone 2
Voeding	**Insecten, nectar en kant-en-klaar voer**
Gehouden	**Solitair of in paren**

KWEEK

Voortplanting	**Eierleggend**
Geslachtsrijp	**4-8 maanden**
Legselgrootte	**2 eieren**
Incubatie	**60-120 dagen** bij 24-28 °C

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
EU-bijlage	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM

The care of *Lygodactylus conraui* is based on the closely related *Lygodactylus williamsi* and on the species' way of life in the wild.

TEMPERATUUR

L. conraui lives in the warm, humid lowlands. By day it should be 24-27 °C, with a basking spot of 32-34 °C that the animal can approach and leave again. At the top of the enclosure it may be 27-29 °C and at the bottom 24-27 °C.

LUCHTVOCHTIGHEID EN WATER

The humidity should be 60-80 %, highest at the bottom.

VERLICHTING EN UVB

The species is diurnal and needs strong, full-spectrum light. Use T5 fluorescent tubes, one with daylight of about 6500 K and one with UVB.

VOEDING

In the wild the species feeds on very small arthropods, especially small moths and termites.

GEDRAG EN OMGANG

Lygodactylus conraui is an animal to watch, not to take in the hand. Dwarf geckos are small, fast and fragile, and handling stresses them and can cost them the tail.

NATUURLIJK VERSPREIDINGSGEBIED

Sierra Leone, Liberia, Ivoorkust, Ghana, Togo, Benin, Nigeria, Kameroen en Equatoriaal-Guinea.