herpy

GEKKO'S

Rüppell's semaphore gecko

Pristurus flavipunctatus

A small, diurnal gecko from the countries around the southern Red Sea, with a tall, serrated tail and yellow spots. It grows to about 8 cm in total length including the tail, and almost nothing is known about its care.

Familie: Sphaerodactylidae Vroegere naam: *Pristurus percristatus; Pristurus gasperetti*

KERNGEGEVENS

HET DIER

Volwassen grootte	**7-8 cm** totale lengte incl. staart
Levensduur	**5-8 jaar**
Actief	**Overdag**
Levenswijze	**Rotsbewonend**
Leefgebied	**Halfwoestijn en rotsachtig terrein**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**40 × 30 × 30 cm** L × B × H voor 1 volwassen dier
Temperatuur overdag	**28-30 °C**
Zonneplek	**40-45 °C**
Temperatuur 's nachts	**20-22 °C**
Luchtvochtigheid	**30-40%**
Water	**Waterbak**
UVB	**Onmisbaar** Fergusonzone 3
Voeding	**Insecten en kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**Solitair of in paren**

KWEEK

Voortplanting	**Eierleggend**
Legselgrootte	**1 eieren**
Incubatie	**70-90 dagen** bij 28 °C

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
EU-bijlage	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM

Almost nothing is known about the care of *Pristurus flavipunctatus*. The care advice is based on the climate of its range and on *Pristurus carteri*, the best-known species of the genus in captivity.

TEMPERATUUR

Create a gradient with 28-30 °C at the cool end and a basking spot of 40-45 °C under a spotlight. At night the temperature may fall to 20-22 °C.

LUCHTVOCHTIGHEID EN WATER

The air must be dry, about 30-40 %. Offer a very shallow dish of fresh water, or put stones in the dish so the animal cannot drown, and mist the enclosure lightly once a week so the animals can drink droplets.

VERLICHTING EN UVB

P. flavipunctatus is diurnal and needs strong daylight and UVB over the basking spot, e.g. a T5 tube. Ferguson zone 3 is an estimate for a diurnal desert gecko.

VOEDING

The species eats small arthropods. Offer springtails, fruit flies, small crickets and other small insects, no wider than the animal's head.

GEDRAG EN OMGANG

Pristurus flavipunctatus is fast and wary and is not suited to handling, and the tail can break off. The behaviour of the animals towards each other in captivity is not documented.

NATUURLIJK VERSPREIDINGSGEBIED

Eritrea, Ethiopië, Djibouti, Somalië, Soedan, Zuid-Soedan, Egypte en Saoedi-Arabië.