herpy

ECHTE HAGEDISSEN

Fringe-fingered Lizard

Acanthodactylus erythrurus

Acanthodactylus erythrurus is a fast, slender lizard from Iberia and north-western Africa with long, fringed toes, brown and grey markings and a reddish tail in juveniles and in females during the breeding season. It reaches 18-23 cm, feeds on insects and needs a deep sand substrate, strong sunlight and high heat.

Familie: Lacertidae

KERNGEGEVENS

HET DIER

Volwassen grootte	**18-23 cm** totale lengte incl. staart
Levensduur	**5-8 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend en gravend**
Leefgebied	**Halfwoestijn en gematigd bos en grasland**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**100 × 50 × 40 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**25-35 °C**
Zonneplek	**40-45 °C**
Temperatuur ’s nachts	**15-22 °C**
Luchtvochtigheid	**30-50%**
Water	**Waterbak**
UVB	**Aanbevolen** Fergusonzone 3
Winterrust	**10-15 °C** 8-12 weken
Voeding	**Insecten**
Gehouden	**In paren**

KWEEK

Voortplanting	**Eierleggend**
Geslachtsrijp	**10-18 maanden**
Legselgrootte	**1-8 eieren**
Legsels per jaar	**1-2**
Incubatie	**35-60 dagen** bij 25-30 °C

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
EU-bijlage	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	Little is known about the care of *Acanthodactylus erythrurus*.
TEMPERATUUR	*Acanthodactylus erythrurus* is very fond of heat: active adults in the wild maintain a body temperature of 32-38 °C.
LUCHTVOCHTIGHEID EN WATER	Humidity should be 30-50 %, as in the dry, open habitat. The sand should be dry, and only the sand in the laying box is kept damp, while the rest of the enclosure is kept well ventilated.
VERLICHTING EN UVB	*Acanthodactylus erythrurus* is diurnal and basks a great deal, so the enclosure needs strong light and UVB.
VOEDING	*Acanthodactylus erythrurus* eats insects and other small arthropods, in the wild especially ants.
GEDRAG EN OMGANG	*Acanthodactylus erythrurus* is fast, wary and shy. It buries itself in the sand when in danger and is hard to catch. Avoid grabbing the animal; the tail is shed under stress. Males are territorial and fight over space.
NATUURLIJK VERSPREIDINGSGEBIED	Spanje, Portugal, Gibraltar, Marokko en Algerije.