herpy

ECHTE HAGEDISSEN

Asian Grass Lizard

Takydromus sexlineatus

Takydromus sexlineatus is a slender, diurnal lizard with an exceptionally long tail, found in tall grass and low vegetation in South-East Asia. It reaches 20-30 cm, most of which is tail, and feeds on small insects. The species is tropical and does not need brumation, but it dries out quickly and needs a tall, densely planted enclosure with moderate humidity.

Familie: Lacertidae

KERNGEGEVENS

HET DIER

Volwassen grootte	**20-30 cm** totale lengte incl. staart
Levensduur	**5-7 jaar**
Actief	**Overdag**
Levenswijze	**Boombewonend en bodembewonend**
Leefgebied	**Savanne**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**60 × 40 × 60 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**25-30 °C**
Zonneplek	**34-40 °C**
Temperatuur 's nachts	**20-23 °C**
Luchtvochtigheid	**60-70%**
Water	**Waterbak**
UVB	**Aanbevolen** Fergusonzone 2
Voeding	**Insecten**
Gehouden	**In paren of één mannetje met meerdere vrouwtjes**

KWEEK

Voortplanting	**Eierleggend**
Legselgrootte	**1-4 eieren**
Legsels per jaar	**2-5**
Incubatie	**40 dagen** bij 26 °C

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
EU-bijlage	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	A small glass tank is too small for *Takydromus sexlineatus*, even though the animal is small. A pair needs at least 60 × 40 × 60 cm (length × width × height), and one male with two females at least 90 × 60 × 60 cm.
TEMPERATUUR	The gradient is the most important thing. The daytime temperature is 25-30 °C, and the surface at the basking spot is 34-40 °C, measured with an infrared thermometer.
LUCHTVOCHTIGHEID EN WATER	Humidity should be 60-70 %. Mist the enclosure with water twice a day so that the lizards can drink droplets from the leaves and walls and so that humidity is kept up.
VERLICHTING EN UVB	*Takydromus sexlineatus* is diurnal. The enclosure needs a small basking lamp and UVB lighting.
VOEDING	*Takydromus sexlineatus* is an insectivore and takes small live insects. Suitable foods are small crickets, flies, fruit flies, springtails, mealworms, waxworms, small locusts and small cockroaches.
GEDRAG EN OMGANG	*Takydromus sexlineatus* is fast, shy and an animal for observation.
NATUURLIJK VERSPREIDINGSGEBIED	India, China, Myanmar (Birma), Thailand, Laos, Cambodja, Vietnam, Maleisië en Indonesië.