herpy

HAZELWORMACHTIGEN

Fringed Arboreal Alligator Lizard

Abronia fimbriata

Abronia fimbriata is a brown, arboreal alligator lizard with thorn-like scales above the ear, found in the cloud forests of Guatemala's north-eastern highlands. It reaches about 25-35 cm, gives birth to live young and eats insects. It needs a tall, airy enclosure with moss and bromeliads and cool temperatures. The species is threatened, and its care is only sparsely documented.

Familie: Anguidae Vroegere naam: *Barissia fimbriata; Gerrhonotus fimbriatus*

KERNGEGEVENS

HET DIER

Volwassen grootte	**25-35 cm** totale lengte incl. staart
Levensduur	**10-20 jaar**
Actief	**Overdag**
Levenswijze	**Boombewonend**
Leefgebied	**Bergbos**
Moeilijkheidsgraad	**Ervaren**

VERZORGING

Min. terrarium	**50 × 50 × 75 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**20-27 °C**
Zonneplek	**26-30 °C**
Temperatuur ’s nachts	**10-18 °C**
Luchtvochtigheid	**60-80%**
Water	**Geen waterbak**
UVB	**Aanbevolen** Fergusonzone 2
Voeding	**Insecten en wormen/slakken**
Gehouden	**Solitair of in paren**

KWEEK

Voortplanting	**Levendbarend**
Geslachtsrijp	**24-36 maanden**
Worpgrootte	**4 jongen**
Legsels per jaar	**1**
Draagtijd	**180-240 dagen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage I**
EU-bijlage	**Bijlage A**
IUCN	**Bedreigd (EN)**

Verzorging in het kort

HET TERRARIUM

Care of *Abronia fimbriata* is only sparsely documented, and the figures given here are borrowed from the closely related *Abronia graminea*, which is the best-known species in the genus.

TEMPERATUUR

Abronia fimbriata does not tolerate heat. The daytime temperature should be 20-27 °C, and the basking spot under a weak spotlight may reach 26-30 °C. The animal is most active at 24-27 °C.

LUCHTVOCHTIGHEID EN WATER

Relative humidity should be 60-80 %, with the highest values at night and after misting. Mist the plants, walls and animals lightly almost every day, ideally with a pressure sprayer or a fogger that produces a fine mist.

VERLICHTING EN UVB

The species is diurnal and needs UVB. The best option is natural, unfiltered sunlight, for example in an outdoor mesh cage in dry weather in summer, with shade and a cool surface so that the animal can move away.

VOEDING

Abronia fimbriata feeds on insects and other small animals. Offer crickets, grasshoppers, mealworms, cockroaches, snails, spiders and soft larvae, for example green caterpillars and fly larvae with a high calcium content.

GEDRAG EN OMGANG

The species is quiet and shy. The animal spends most of the day sitting still on a branch or in a bromeliad and basks for short periods. It is not an animal to be handled.

NATUURLIJK VERSPREIDINGSGEBIED

Guatemala.