herpy

KAMELEONS

Von Hoehnel's Chameleon

Trioceros hoehnelii

Trioceros hoehnelii is a small, live-bearing chameleon from the cool highlands of Kenya and Uganda, where nights can approach freezing. The male has a tall casque and a single horn on the snout. The species is robust against cold, but tolerates heat badly, so it needs a cool terrarium with a clear drop in temperature at night and high night-time humidity.

Familie: Chamaeleonidae Vroegere naam: *Chamaeleo hoehnelii; Chamaeleo bergeri; Chamaeleo leikipiensis*

KERNGEGEVENS

HET DIER

Volwassen grootte	**18-25 cm** totale lengte incl. staart
Gewicht	**15-70 g**
Levensduur	**5-6 jaar**
Actief	**Overdag**
Levenswijze	**Boombewonend**
Leefgebied	**Bergbos**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**60 × 45 × 90 cm** L × B × H voor 1 volwassen dier
Temperatuur overdag	**20-22 °C**
Zonneplek	**24-25 °C**
Temperatuur ’s nachts	**8-14 °C**
Luchtvochtigheid	**50-100%**
UVB	**Onmisbaar** Fergusonzone 3
Voeding	**Insecten**
Gehouden	**Solitair**

KWEEK

Voortplanting	**Levendbarend**
Geslachtsrijp	**12 maanden**
Worpgrootte	**4-22 jongen**
Legsels per jaar	**1-2**
Draagtijd	**120-180 dagen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
EU-bijlage	**Bijlage B**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM

An adult needs a terrarium of at least 60 × 45 × 90 cm (length × width × height). The size is borrowed from similar small mountain chameleons of the genus *Trioceros* and is a minimum.

TEMPERATUUR

The species comes from cool highlands and tolerates heat badly. The daytime temperature in the terrarium should be 20-22 °C with a basking spot of 24-25 °C. Young animals can die if the temperature goes higher.

LUCHTVOCHTIGHEID EN WATER

Night is the species' humid time. The humidity should rise towards 100 % at night, with mist from an ultrasonic fogger for about three hours late at night and early in the morning. By day the humidity may fall to about 50-80 %.

VERLICHTING EN UVB

The species is diurnal and uses a lot of light. Give 12 hours of light a day all year. A T5 HO tube of 6500 K or a powerful LED lamp across the width of the terrarium provides the general light. UVB is necessary.

VOEDING

T. hoehnelii is insectivorous and takes most insects of suitable size. Offer crickets, small grasshoppers, flies, cockroach nymphs and other small insects, and keep the prey to no more than the distance between the animal's eyes.

GEDRAG EN OMGANG

The species is kept singly. Males are very aggressive towards each other and towards similar species: they hiss, puff themselves up and turn black, so males must not be able to see each other.

NATUURLIJK VERSPREIDINGSGEBIED

Kenia en Oeganda.