herpy

KIKKERS

Large-webbed Bell Toad

Bombina maxima

Bombina maxima is the largest species in the genus: a stout, grey-brown, heavily warty frog of 6-8 cm with an orange belly, marbled with black, which it displays when it feels threatened. It comes from the mountains of south-western China and wants clean, cool water and a cool terrarium. It is easy to keep when the temperature is held down, but makes greater demands on water quality than *Bombina orientalis*.

Familie: Bombinatoridae Vroegere naam: *Bombinator maximus* **Mild huidgif**

KERNGEGEVENS

HET DIER

Volwassen grootte	**6-8 cm** kop-romplengte
Levensduur	**6-15 jaar**
Actief	**Overdag en in de schemering**
Levenswijze	**Semi-aquatisch**
Leefgebied	**Moerasgebied en gematigd bos en grasland**
Huidgif	**Mild** Was je handen na contact en houd je vingers uit de buurt van je ogen en mond
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**80 × 40 × 40 cm** L × B × H voor 4 volwassen dieren
Temperatuur overdag	**18-23 °C**
Temperatuur 's nachts	**12-17 °C**
Luchtvochtigheid	**60-85%**
Water	**Aquaterrarium** diepte 5-10 cm
UVB	**Optioneel**
Winterrust	**4-8 °C** 8-12 weken
Voeding	**Insecten en wormen/slakken**
Gehouden	**In groepen**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes in het water**
Legplaats	**In het water**
Geslachtsrijp	**24-36 maanden**
Eieren per legsel	**20-150**
Uitkomen	**3-10 dagen** bij 18-22 °C
Larvenperiode	**30-90 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Geen**

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	The species needs an aquaterrarium in which about two thirds of the floor area is water and one third land. Four adults need at least 80 × 40 × 40 cm.
TEMPERATUUR	The species comes from cool highlands and must be kept cooler than most frogs. By day 18-23 °C is suitable, and the temperature should never exceed 25 °C. At night it may fall to 12-17 °C.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 60-85 %. The large water section does most of the work, so it is enough to mist the land section lightly once a day. Moss and soil on the land section must be damp but not wet, and there must be damp hides.
WATER EN WATERKWALITEIT	The water must be 5-10 cm deep, and the animals must always be able to climb out easily, for example along sloping roots, flat stones or plants. The water is at room temperature and should not be heated.
VERLICHTING EN UVB	Give 12 hours of light a day in the summer half of the year and slightly shorter days in winter, controlled by a timer. An LED fixture with a daylight spectrum gives enough light for the plants.
VOEDING	The food must move. Give ordinary feeder insects of a suitable size: crickets, flies, woodlice, earthworms and small snails. Mealworms are fatty and are given only once in a while.
GEDRAG EN OMGANG	The species is active by day and at dusk and is often seen out in the open, on land or floating at the surface. It is best kept in a group. In the breeding season things get lively when the males call from the water.
NATUURLIJK VERSPREIDINGSGEBIED	China.