herpy

KIKKERS

Yellow-bellied Toad

Bombina variegata

Bombina variegata is a small, compact frog of 3-6 cm with a grey-brown, sharp-warted back and a bright yellow and blue-grey belly, which it displays when it feels threatened. The species lives in the mountains and hill country of central and southern Europe, favours small, sunny waterholes and is hardy, inquisitive and long-lived in the terrarium. It needs cool temperatures and a cool hibernation, but otherwise little space.

Familie: Bombinatoridae **Mild huidgif**

KERNGEGEVENS

HET DIER

Volwassen grootte	**3-6 cm** kop-romplengte
Gewicht	**2,3-12 g**
Levensduur	**10-20 jaar**
Actief	**In de schemering en 's nachts**
Levenswijze	**Semi-aquatisch**
Leefgebied	**Moerasgebied en gematigd bos en grasland**
Huidgif	**Mild** Was je handen na contact en houd je vingers uit de buurt van je ogen en mond
Moeilijkheidsgraad	**Beginner**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes in het water**
Legplaats	**In het water**
Geslachtsrijp	**12-24 maanden**
Eieren per legsel	**40-170**
Legsels per jaar	**1-3**
Uitkomen	**2-3 dagen**
Larvenperiode	**25-70 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Geen**

VERZORGING

Min. terrarium	**60 × 30 × 30 cm** L × B × H voor 4 volwassen dieren
Temperatuur overdag	**20-24 °C**
Temperatuur 's nachts	**16-20 °C**
Luchtvochtigheid	**60-80%**
Water	**Watergedeelte** diepte 5-10 cm
UVB	**Optioneel**
Winterrust	**5-10 °C** 8-12 weken
Voeding	**Insecten en wormen/slakken**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM

There are not many detailed husbandry accounts of *Bombina variegata*. The recommendations here are based on the species' way of life and on the close relative *Bombina orientalis*, which is far better known in captivity.

TEMPERATUUR

The species is adapted to a temperate climate and should not be given heat like most terrarium animals. By day 20-24 °C is suitable, and at night it should be slightly cooler, 16-20 °C.

LUCHTVOCHTIGHEID EN SPROEIEN

Humidity should be 60-80 %. The large water section holds most of the humidity, so it is normally enough to mist the land section lightly once a day, especially in a dry room in winter.

WATER EN WATERKWALITEIT

The water section should fill about half the floor area and be 5-10 cm deep.

VERLICHTING EN UVB

Give 12-14 hours of light in summer and down to about 10 hours in winter, controlled by a timer. An LED fixture provides the basic light and makes the plants thrive. No heat lamp is needed. UVB is optional.

VOEDING

B. variegata has a large appetite and takes anything live that is small enough to fit in the mouth: crickets, small crickets, fruit flies, flies, woodlice, silverfish, small earthworms and small snails.

GEDRAG EN OMGANG

B. variegata is inquisitive and sociable and can be kept in pairs or in groups. The animals should be roughly the same size, because large animals can eat small ones, and at feeding they can become a little pushy.

NATUURLIJK VERSPREIDINGSGEBIED

Frankrijk, België, Nederland, Luxemburg, Duitsland, Zwitserland, Liechtenstein, Oostenrijk, Italië, Tsjechië, Slowakije, Polen, Hongarije, Roemenië, Moldavië, Oekraïne, Slovenië, Kroatië, Bosnië en Herzegovina, Servië, Montenegro, Noord-Macedonië, Albanië, Bulgarije en Griekenland.