herpy

KIKKERS

Edible Bullfrog

Pyxicephalus edulis

Pyxicephalus edulis is a stout, wide-mouthed burrowing frog from Africa's savannas and flooded grasslands: smaller than *Pyxicephalus adspersus*, but with the same appetite, the same tooth-like jaw projections and a bite that must be taken seriously. It spends most of its time buried, is robust and can be kept in a relatively small terrarium.

Familie: Pyxicephalidae

KERNGEGEVENS

HET DIER

Volwassen grootte	**8-12 cm** kop-romplengte
Levensduur	**10-15 jaar**
Actief	**In de schemering en 's nachts**
Levenswijze	**Bodembewonend en gravend**
Leefgebied	**Savanne en moerasgebied**
Huidgif	**Geen bekend**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**60 × 45 × 30 cm** L × B × H voor 1 volwassen dier
Temperatuur overdag	**23-26 °C**
Temperatuur 's nachts	**18-22 °C**
Luchtvochtigheid	**50-70%**
Water	**Waterbak**
UVB	**Optioneel** Fergusonzone 1
Voeding	**Insecten, gewervelde dieren en wormen/slakken**
Gehouden	**Solitair**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes in het water**
Legplaats	**In het water**
Eieren per legsel	**3.000-4.000**
Uitkomen	**2-3 dagen** bij 24-27 °C
Larvenperiode	**14-35 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren**

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	Little is known about the care of *Pyxicephalus edulis*, and most of what is written about bullfrogs concerns the larger *Pyxicephalus adspersus*.
TEMPERATUUR	By day the terrarium should hold 23-26 °C and at night 18-22 °C. The temperature should be steady and must not exceed 28 °C for any length of time: even a rise from 25 to 30 °C raises stress hormones, and the animals lose weight.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 50-70 %. It comes from the moist substrate and the water dish, and the terrarium is misted daily with water without chlorine, so that the substrate stays slightly moist.
WATER EN WATERKWALITEIT	The frog must always have access to a large, flat dish in which the whole animal can lie and bathe.
VERLICHTING EN UVB	The species is most active at dusk and at night and does not need strong light.
VOEDING	*Pyxicephalus edulis* is a predator that eats almost anything it can overpower. In the wild the diet consists especially of beetles and grasshoppers, as well as termites, millipedes, ants and spiders.
GEDRAG EN OMGANG	The animal almost always lies buried with only the eyes and nostrils above the substrate, waiting for prey. It is most active at dusk and at night and comes out to hunt and swim.
NATUURLIJK VERSPREIDINGSGEBIED	Mauritanië, Mali, Burkina Faso, Benin, Nigeria, Tsjaad, Kameroen, Angola, Zambia, Kenia, Tanzania, Malawi, Mozambique, Zimbabwe, Botswana en Zuid-Afrika.