herpy

KIKKERS

Black-webbed Treefrog

Rhacophorus kio

Rhacophorus kio is a large, grass-green flying frog from the rainforests of southern China, Laos, Thailand and Vietnam. Females reach up to 8 cm. The webbing is orange-yellow with a black spot at the base of each finger and toe, and the frog spreads it out when it glides between the trees. The species is nocturnal and shy, and little is known about its care in captivity.

Familie: Rhacophoridae

KERNGEGEVENS

HET DIER

Volwassen grootte	**6-8 cm** kop-romplengte
Levensduur	**5-10 jaar**
Actief	**'s nachts**
Levenswijze	**Boombewonend**
Leefgebied	**Regenwoud en bergbos**
Huidgif	**Geen bekend**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**60 × 45 × 90 cm** L × B × H voor 3 volwassen dieren
Temperatuur overdag	**21-24 °C**
Temperatuur 's nachts	**18-21 °C**
Luchtvochtigheid	**70-90%**
Water	**Waterbak** diepte 2-3 cm
UVB	**Aanbevolen** Fergusonzone 1
Voeding	**Insecten**
Gehouden	**Solitair, in paren of in groepen**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes in het water**
Legplaats	**In een schuimnest**
Eieren per legsel	**130-180**
Uitkomen	**5 dagen**
Larvenperiode	**45-85 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Geen**

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	Little is known about keeping *Rhacophorus kio*.
TEMPERATUUR	Keep the terrarium at 21-24 °C by day, corresponding to ordinary room temperature, and let it fall to 18-21 °C at night. The frog tolerates short periods up to 27-28 °C, but the temperature must not exceed 28 °C.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 70-90 %, lowest by day and highest at night.
WATER EN WATERKWALITEIT	Place a stable, shallow water bowl in the terrarium, which the frog can sit in at night. 2-3 cm of water is enough, so that the frog easily gets out again.
VERLICHTING EN UVB	The frog is arboreal, and a weak UVB source of the type used for tropical forest animals is recommended. A low UV index on the uppermost leaves (Ferguson zone 1) is a cautious starting point.
VOEDING	*Rhacophorus kio* eats live insects. Crickets can be used throughout its life, from small newly hatched ones to large ones, supplemented with small cockroaches, soldier fly larvae and an occasional wax moth larva as a treat.
GEDRAG EN OMGANG	The species is nocturnal and shy and spends the day hidden among the leaves; most of the time it sits on the glass or on the furnishings.
NATUURLIJK VERSPREIDINGSGEBIED	China, Laos, Thailand en Vietnam.