herpy

KIKKERS

Tonkin Bug-eyed Frog

Theloderma corticale

Theloderma corticale is a nocturnal, semi-aquatic frog from the forests of northern Vietnam, Laos and southern China. The warty, green and black-blotched skin resembles moss and lichen so well that the frog almost disappears in the enclosure. It wants cool and wet conditions and does not tolerate heat, but is on the other hand easy to get to breed.

Familie: Rhacophoridae Vroegere naam: *Rhacophorus corticalis*

KERNGEGEVENS

HET DIER

Volwassen grootte	**6-9 cm** kop-romplengte
Levensduur	**10-20 jaar**
Actief	**'s nachts**
Levenswijze	**Boombewonend en semi-aquatisch**
Leefgebied	**Regenwoud, bergbos en rotsachtig terrein**
Huidgif	**Geen bekend**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**80 × 50 × 40 cm** L × B × H voor 4 volwassen dieren
Temperatuur overdag	**18-23 °C**
Temperatuur 's nachts	**15-20 °C**
Luchtvochtigheid	**80-90%**
Water	**Aquaterrarium** diepte 5-10 cm
Watertemperatuur	**17-22 °C**
UVB	**Optioneel** Fergusonzone 1
Winterrust	**13-16 °C** 4-22 weken
Voeding	**Insecten**
Gehouden	**In paren of in groepen**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes in het water**
Legplaats	**Op bladeren boven het water**
Geslachtsrijp	**8-10 maanden**
Eieren per legsel	**6-47**
Uitkomen	**9-22 dagen**
Larvenperiode	**75-180 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Kant-en-klaar voer**
Broedzorg	**Geen**

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	*Theloderma corticale* needs an aquaterrarium where a large part of the floor, or the whole floor, stands under water.
TEMPERATUUR	Heat is the species' greatest enemy. By day there should be 18-23 °C and at night 15-20 °C. Down to about 10 °C is fine, while anything above 25 °C is a warning sign, and temperatures above 27 °C can kill the frogs.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity must be high: over 80% and never below 60%, otherwise the frogs may dry out. The females lay eggs at 80-90%. The large water surface keeps humidity up by itself, so the frogs do not need misting very often.
WATER EN WATERKWALITEIT	The water section is the species' home. The water should be 5-10 cm deep, and deeper water also works. The frogs must be able to sit in the water and easily climb onto cork or roots.
VERLICHTING EN UVB	The frog is nocturnal and has no documented need for UVB. A low-output UVB tube for shade-dwelling species (Ferguson zone 1) can be used and provides the small UV index that can support the day-night rhythm, but it is optional.
VOEDING	*T. corticale* eats insects: well gut-loaded crickets, dubia roaches, waxworms, black soldier fly larvae and small hornworms.
GEDRAG EN OMGANG	*Theloderma corticale* is calm and hides by day in tubes, caves and among plants. A few minutes after the light is switched off, it becomes active.
NATUURLIJK VERSPREIDINGSGEBIED	Vietnam, Laos en China.