herpy

KIKKERS

Chantaburi Bug-eyed Frog

Theloderma stellatum

Theloderma stellatum is a small, tree-dwelling mossy frog from the lowland rainforest of Thailand, Cambodia, Laos and Vietnam. It resembles a darker version of *Theloderma asperum*, grows to about 3-4 cm and lays its eggs above water in water-filled tree hollows. Its care is similar to that of its relatives, but almost all figures are borrowed from them, because the species is rare in captivity.

Familie: Rhacophoridae

KERNGEGEVENS

HET DIER

Volwassen grootte	**3-4 cm** kop-romplengte
Levensduur	**5-10 jaar**
Actief	**'s nachts**
Levenswijze	**Boombewonend en semi-aquatisch**
Leefgebied	**Regenwoud**
Huidgif	**Geen bekend**
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes in het water**
Legplaats	**Op bladeren boven het water**
Geslachtsrijp	**8-10 maanden**
Eieren per legsel	**4-15**
Uitkomen	**10-14 dagen**
Larvenperiode	**75-180 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Kant-en-klaar voer**
Broedzorg	**Geen**

VERZORGING

Min. terrarium	**45 × 45 × 45 cm** L × B × H voor 4 volwassen dieren
Temperatuur overdag	**20-24 °C**
Temperatuur 's nachts	**17-21 °C**
Luchtvochtigheid	**75-95%**
Water	**Watergedeelte** diepte 5-10 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Insecten**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	Little documented knowledge exists on the care of *Theloderma stellatum*.
TEMPERATUUR	By day the temperature should be 20-24 °C and at night 17-21 °C.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 75-95 % and never below 60 %, otherwise the frogs may dry out. The large water surface keeps the humidity up on its own, so the frogs do not need much misting.
WATER EN WATERKWALITEIT	The frog spends a lot of time at the water's edge and even sits under water, so the water section is not there for show.
VERLICHTING EN UVB	The frog is nocturnal and has no documented need for UVB. A weak UVB tube for shade species (Ferguson zone 1) can be used and is optional, but the frogs must always be able to retreat into shade.
VOEDING	*T. stellatum* eats insects. Small crickets are the main food, supplemented with waxworms, flies, springtails for the smallest young, small dubia roaches and soldier fly larvae.
GEDRAG EN OMGANG	*Theloderma stellatum* is nocturnal and sits hidden by day in tubes, among plants or close to the water. The frog does well in a group as long as there is enough space. The males call at night.
NATUURLIJK VERSPREIDINGSGEBIED	Thailand, Cambodja, Laos en Vietnam.