herpy

KIKKERS

Amazon Milk Frog

Trachycephalus resinifictrix

Trachycephalus resinifictrix is a large, powerful, nocturnal tree frog from the Amazon canopy. The juveniles are striped black and white, the adults marbled, and when threatened the frog secretes a milky-white skin secretion. It breeds in water-filled tree holes, is robust to keep and in many ways resembles *Ranoidea caerulea*.

Familie: Hylidae Vroegere naam: *Phrynohyas resinifictrix* **Mild huidgif**

KERNGEGEVENS

HET DIER

Volwassen grootte	**6-10 cm** kop-romplengte
Levensduur	**8-10 jaar**
Actief	**'s nachts**
Levenswijze	**Boombewonend**
Leefgebied	**Regenwoud**
Huidgif	**Mild** Was je handen na contact en houd je vingers uit de buurt van je ogen en mond
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes in het water**
Legplaats	**In het water**
Geslachtsrijp	**12-18 maanden**
Eieren per legsel	**100-1.500**
Uitkomen	**1-2 dagen**
Larvenperiode	**21-75 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Kant-en-klaar voer**
Broedzorg	**Geen**

VERZORGING

Min. terrarium	**75 × 30 × 45 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**21-29 °C**
Temperatuur 's nachts	**18-21 °C**
Luchtvochtigheid	**50-100%**
Water	**Waterbak**
UVB	**Aanbevolen** Fergusonzone 2
Voeding	**Insecten en wormen/slakken**
Gehouden	**Solitair, in paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	An adult pair needs an enclosure of at least 75 × 30 × 45 cm (length × width × height), and that is enough for up to four adults. A group of five needs at least 60 × 45 × 60 cm.
TEMPERATUUR	*Trachycephalus resinifictrix* tolerates a wide temperature range.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity is less critical than temperature, but the enclosure should be misted at least once a day. Use dechlorinated tap water or reverse-osmosis water, which also leaves no limescale marks on the glass.
WATER EN WATERKWALITEIT	*Trachycephalus resinifictrix* needs a large bowl of clean water, corresponding to the tree hole in the wild. The bowl must be large enough for the frog to sit in, and it is used almost every night, so do not skimp on size.
VERLICHTING EN UVB	*Trachycephalus resinifictrix* is nocturnal and can both live and breed without UVB, but juveniles raised with UVB grow larger and develop a healthier skeleton than juveniles without.
VOEDING	*T. resinifictrix* is greedy and rarely fussy. Crickets can make up most of the diet, and they should be roughly as long as the frog's head is wide.
GEDRAG EN OMGANG	*T. resinifictrix* is not an animal for handling. The skin is permeable and sensitive to oils and salts from human skin, and the skin secretion can be irritating.
NATUURLIJK VERSPREIDINGSGEBIED	Brazilië, Frans-Guyana, Suriname, Guyana, Venezuela, Colombia, Peru en Ecuador.

De volledige gids met kaart en updates: herpy.org/en/frogs/trachycephalus-resinifictrix