herpy

KIKKERS

Milk Frog

Trachycephalus typhonius

Trachycephalus typhonius is a large, nocturnal tree frog with warty skin and golden eye rims. It lives over an enormous area from Central America to northern Argentina, from rainforest to savanna and gardens. The skin secretes a white substance that stings the mucous membranes, and experience of keeping it is limited.

Familie: Hylidae Vroegere naam: *Phrynohyas venulosa* **Mild huidgif**

KERNGEGEVENS

HET DIER

Volwassen grootte	**6-8 cm** kop-romplengte
Levensduur	**8-10 jaar**
Actief	**'s nachts**
Levenswijze	**Boombewonend**
Leefgebied	**Savanne, droog bos en regenwoud**
Huidgif	**Mild** Was je handen na contact en houd je vingers uit de buurt van je ogen en mond
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes in het water**
Legplaats	**In het water**
Eieren per legsel	**100-1.500**
Uitkomen	**1-2 dagen**
Larvenperiode	**21-75 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Geen**

VERZORGING

Min. terrarium	**60 × 40 × 60 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**24-28 °C**
Temperatuur 's nachts	**18-22 °C**
Luchtvochtigheid	**60-90%**
Water	**Waterbak**
UVB	**Aanbevolen** Fergusonzone 2
Voeding	**Insecten**
Gehouden	**Solitair, in paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM
The care of *Trachycephalus typhonius* is based on the closely related *Trachycephalus resinifictrix*, which is a similarly large, nocturnal tree frog.

TEMPERATUUR
Keep the enclosure at 24-28 °C by day and 18-22 °C at night. There is no winter rest period. The frog tolerates a brief deviation, but the temperature must not stay above 30 °C for long.

LUCHTVOCHTIGHEID EN SPROEIEN
Humidity should be 60-90 % with a rise after misting. The species also lives in dry habitats and is hardy, so the air does not need to be saturated all the time.

WATER EN WATERKWALITEIT
Put a large, stable water bowl in the enclosure. The bowl must be large enough for the frog to sit in, and the water must be shallow, so that the animal can easily get out again.

VERLICHTING EN UVB
Provide 10-12 hours of light a day, first and foremost for the plants. The frog sleeps by day and is most active in the dark, so it does not need strong light. A low level of UVB helps the frog absorb the calcium in the food.

VOEDING
The frog takes most things it can manage: crickets, houseflies, blowflies, cockroaches, worms and other insects. Crickets can make up the bulk of the diet, and the other feeder insects are given in rotation.

GEDRAG EN OMGANG
Trachycephalus typhonius is nocturnal and sleeps by day in hiding, often high up, and becomes active when the light is switched off.

NATUURLIJK VERSPREIDINGSGEBIED
Mexico, Belize, Guatemala, Honduras, El Salvador, Nicaragua, Costa Rica, Panama, Colombia, Venezuela, Trinidad en Tobago, Guyana, Suriname, Frans-Guyana, Brazilië, Bolivia, Paraguay en Argentinië.