herpy

LEGUANEN

Common Puerto Rican Anole

Anolis cristatellus

Anolis cristatellus is a stocky, diurnal anole from Puerto Rico and the Virgin Islands. The male has a tall, fin-like crest on the tail that is always erect, and a greenish-yellow dewlap with an orange-red margin, which he extends while bobbing and doing push-ups. The species lives on trunks and low down in the vegetation, is territorial and very shy, and needs UVB, live food and an enclosure with both branches and floor space.

Familie: Dactyloidae Vroegere naam: *Anolis cozumelae; Anolis lindeni; Ctenonotus cristatellus*

KERNGEGEVENS

HET DIER

Volwassen grootte	**15-20 cm** totale lengte incl. staart
Levensduur	**4-8 jaar**
Actief	**Overdag**
Levenswijze	**Boombewonend en bodembewonend**
Leefgebied	**Droog bos en regenwoud**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**60 × 45 × 60 cm** L × B × H voor 1 volwassen dier
Temperatuur overdag	**26-29 °C**
Zonneplek	**32-35 °C**
Temperatuur ’s nachts	**20-24 °C**
Luchtvochtigheid	**60-80%**
Water	**Waterbak**
UVB	**Onmisbaar** Fergusonzone 2
Voeding	**Insecten**
Gehouden	**Solitair of één mannetje met meerdere vrouwtjes**

KWEEK

Voortplanting	**Eierleggend**
Legselgrootte	**1-2 eieren**
Incubatie	**45-55 dagen** bij 26-28 °C

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
EU-bijlage	**Niet opgenomen**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	Little is known about keeping *Anolis cristatellus*.
TEMPERATUUR	The air should be 26-29 °C by day, and the basking spot should have a surface temperature of 32-35 °C, measured with an infrared thermometer. The cooler side is 24-26 °C. At night 20-24 °C is suitable.
LUCHTVOCHTIGHEID EN WATER	Humidity should be 60-80 %: lower in the middle of the day and higher at night.
VERLICHTING EN UVB	*Anolis cristatellus* is diurnal and needs UVB every day to produce vitamin D3 and absorb calcium. The UVB lamp should give Ferguson zone 2, i.e. a moderate UV index at the basking spot.
VOEDING	The species is insectivorous and takes live prey that moves: crickets, cockroaches (such as dubia and discoid), grasshoppers, silkworms, hornworms, small superworms and soldier fly larvae.
GEDRAG EN OMGANG	The species is shy, quick and not an animal for handling. It can drop its tail, and it does not grow back in the same form.
NATUURLIJK VERSPREIDINGSGEBIED	Puerto Rico, Amerikaanse Maagdeneilanden en Britse Maagdeneilanden.