herpy

LEGUANEN

Blue Spiny Lizard

Sceloporus cyanogenys

Sceloporus cyanogenys is a stocky, rough-scaled lizard from rocky country in southern Texas and eastern Mexico. The male has a blue throat, blue sides and a black collar. The species is diurnal, feeds on insects and needs a roomy, dry enclosure with rocks, UVB and a hot basking spot. It gives birth to live young and is shy and fast.

Familie: Phrynosomatidae

KERNGEGEVENS

HET DIER

Volwassen grootte	**25-36 cm** totale lengte incl. staart
Levensduur	**5-7 jaar**
Actief	**Overdag**
Levenswijze	**Rotsbewonend en bodembewonend**
Leefgebied	**Halfwoestijn en rotsachtig terrein**
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**100 × 50 × 50 cm** L × B × H voor 1 volwassen dier
Temperatuur overdag	**27-31 °C**
Zonneplek	**38-42 °C**
Temperatuur 's nachts	**21-26 °C**
Luchtvochtigheid	**30-50%**
Water	**Waterbak**
UVB	**Onmisbaar** Fergusonzone 3
Voeding	**Insecten**
Gehouden	**Solitair, in paren of één mannetje met meerdere vrouwtjes**

KWEEK

Voortplanting	**Levendbarend**
Geslachtsrijp	**28-36 maanden**
Worpgrootte	**5-20 jongen**
Legsels per jaar	**1**
Draagtijd	**90-120 dagen**

BESCHERMING EN PAPIEREN

CITES	**Niet opgenomen**
EU-bijlage	**Niet opgenomen**
IUCN	**Niet beoordeeld (NE)**

Verzorging in het kort

HET TERRARIUM	An adult needs an enclosure of at least 100 × 50 × 50 cm (length × width × height).
TEMPERATUUR	The enclosure should have a clear gradient. The air should be 27-31 °C during the day, and the basking spot should be 38-42 °C on the surface, measured with an infrared thermometer.
LUCHTVOCHTIGHEID EN WATER	The air should be dry to moderately humid: 30-50 %. The species does poorly in humid, stagnant air, so ventilation should be good. Provide a low, stable bowl of fresh water, and change the water daily.
VERLICHTING EN UVB	*Sceloporus cyanogenys* is diurnal and basks in full sun, so it needs strong light and UVB. Use a T5 HO tube (Ferguson zone 3) in a reflector over the basking spot, and keep the distance within the manufacturer's recommendation.
VOEDING	*Sceloporus cyanogenys* eats insects. Give a varied diet of live insects: crickets, small cockroaches, mealworms, morio worms, silkworms and waxworms (the last sparingly).
GEDRAG EN OMGANG	The animal is fast and shy and defends itself by biting and fleeing. It is an animal to look at, not to handle. If it has to be picked up, hold the body from neck to vent.
NATUURLIJK VERSPREIDINGSGEBIED	Verenigde Staten, Mexico en Guatemala.