herpy

PIJLGIFKIKKERS

Brazil-nut Poison Frog

Adelphobates castaneoticus

Adelphobates castaneoticus is a small, glossy black poison frog with white or yellowish spots and bright orange patches at the forelimbs and knees. It lives on the forest floor in the Brazilian state of Pará in the Amazon and is named after the Brazil nut, whose empty shells the male uses as water pools for the tadpoles. Little is known about its care.

Familie: Dendrobatidae Vroegere naam: *Dendrobates castaneoticus*

KERNGEGEVENS

HET DIER

Volwassen grootte	**18-23 mm** kop-romplengte
Levensduur	**8-15 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Ervaren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Geslachtsrijp	**5-7 maanden**
Eieren per legsel	**4-8**
Uitkomen	**10-14 dagen**
Larvenperiode	**60-90 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en andere kikkervisjes**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

VERZORGING

Min. terrarium	**45 × 45 × 45 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**24-27 °C**
Temperatuur ’s nachts	**20-24 °C**
Luchtvochtigheid	**70-100%**
Water	**Geen waterbak**
UVB	**Optioneel**
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM

Little is known about the care of *Adelphobates castaneoticus*. The species lives in the leaf litter of the forest floor in a warm, humid lowland climate.

TEMPERATUUR

The daytime temperature should be 24-27 °C, and at night it falls 3-4 °C, to 20-24 °C. The variation over the year should be small, 1-2 °C. This fits the lowland climate of Pará. There should be no basking spot and no heat lamp.

LUCHTVOCHTIGHEID EN SPROEIEN

Humidity should be about 70-80 % in the middle of the day and close to 100 % in the morning and evening, and the leaf litter and substrate should be moist at all times.

WATER EN WATERKWALITEIT

A water dish is not necessary unless you want to raise young. The frogs get their moisture from misting and the moist leaf litter.

VERLICHTING EN UVB

Provide 10-12 hours of light a day with fluorescent tubes or LEDs for the plants. In the wild the species lives in the shade under the forest canopy, so the light should give the plants enough to grow without heating the enclosure.

VOEDING

In the wild *A. castaneoticus* feeds on ants, termites and other small arthropods. In captivity feed springtails, small fruit flies and small isopods.

GEDRAG EN OMGANG

A. castaneoticus is diurnal and lives mostly on the forest floor. Temperament and territorial behaviour are not documented, but *A. galactonotus* can become aggressive towards conspecifics when it becomes sexually mature.

NATUURLIJK VERSPREIDINGSGEBIED

Brazilië.