herpy

PIJLGIFKIKKERS

Rio Madeira Poison Frog

Adelphobates quinquevittatus

Adelphobates quinquevittatus is a very small, glossy black poison frog of 14-16 mm with white stripes and grey, greenish or orange legs. It lives in the leaf litter of the rainforest in the Madeira River drainage in the southern Amazon and is only rarely kept. Very little is known about its care.

Familie: Dendrobatidae Vroegere naam: *Dendrobates quinquevittatus*

KERNGEGEVENS

HET DIER

Volwassen grootte	**14-16 mm** kop-romplengte
Levensduur	**8-15 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Ervaren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Eieren per legsel	**4-8**
Uitkomen	**10-14 dagen**
Larvenperiode	**60-90 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en andere kikkervisjes**
Broedzorg	**Draagt de kikkervisjes**

VERZORGING

Min. terrarium	**45 × 45 × 45 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**24-27 °C**
Temperatuur 's nachts	**20-24 °C**
Luchtvochtigheid	**70-100%**
Water	**Geen waterbak**
UVB	**Optioneel**
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	Very little is known about the care of *Adelphobates quinquevittatus*. The species lives in the leaf litter in a warm and humid lowland climate.
TEMPERATUUR	The daytime temperature should be 24-27 °C, and at night it falls 3-4 °C, to 20-24 °C. The variation over the year should be small, 1-2 °C. This fits the lowland climate of Rondônia.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be about 70-80 % in the middle of the day and close to 100 % in the morning and evening, and the leaf litter and substrate should be moist at all times.
WATER EN WATERKWALITEIT	A water dish is not necessary unless you want to raise young. The frogs get their moisture from misting and the moist leaf litter.
VERLICHTING EN UVB	Provide 10-12 hours of light a day with fluorescent tubes or LEDs for the plants. In the wild the species lives in the shade under the forest canopy, so the light should give the plants enough to grow without heating the enclosure.
VOEDING	In the wild *A. quinquevittatus* feeds mostly on ants and termites. In captivity feed small fruit flies (*Drosophila melanogaster*), springtails, micro crickets, small waxworm larvae, sieved meadow plankton and small isopods.
GEDRAG EN OMGANG	*A. quinquevittatus* is diurnal and lives mostly in the leaf litter. Temperament and territorial behaviour are not documented.
NATUURLIJK VERSPREIDINGSGEBIED	Brazilië.