herpy

PIJLGIFKIKKERS

Brilliant-thighed Poison Frog

Allobates femoralis

Allobates femoralis is a small, dark frog of 3-4 cm from the Amazon and the Guiana Shield with an orange-yellow spot behind the arm and a crescent-shaped orange spot on the thigh. It is diurnal and ground-dwelling, the males defend territories with a loud call, and the courtship is one of the longest among poison frogs. The species belongs to the family Aromobatidae and is regarded as non-toxic.

Familie: Aromobatidae Vroegere naam: *Epipedobates femoralis; Phyllobates femoralis; Dendrobates femoralis*

KERNGEGEVENS

HET DIER

Volwassen grootte	**3-4 cm** kop-romplengte
Levensduur	**5-10 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif	**Geen bekend**
Moeilijkheidsgraad	**Ervaren**

VERZORGING

Min. terrarium	**60 × 40 × 40 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**25-28 °C**
Temperatuur 's nachts	**21-24 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren of één mannetje met meerdere vrouwtjes**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Eieren per legsel	**8-20**
Uitkomen	**15-20 dagen**
Larvenperiode	**40-50 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	A pair of *Allobates femoralis* needs a glass terrarium of at least 60 × 40 × 40 cm. The species lives near the ground, so floor area counts most, but the males like to climb a little to call.
TEMPERATUUR	By day the temperature should be 25-28 °C and at night 21-24 °C, that is 3-4 degrees lower. The daily swing corresponds to conditions in French Guiana and promotes breeding.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be high: 70-80 % in the middle of the day and up to 100 % morning and evening.
WATER EN WATERKWALITEIT	Provide a shallow glass dish of about 10 cm in diameter with 1-2 cm of water, which the frogs can easily climb out of. The male also uses it when he is to deposit the tadpoles.
VERLICHTING EN UVB	The light is mainly for the plants. Provide 12 hours of light a day, for example from 7 to 19 h. Fluorescent tubes or LEDs of around 6,500 K give natural colours and good plant growth.
VOEDING	*Allobates femoralis* feeds on small insects. In the enclosure the staple diet consists of wingless fruit flies, supplemented with springtails, very small crickets, small wax moth larvae and meadow plankton.
GEDRAG EN OMGANG	*Allobates femoralis* is diurnal, and the males are territorial.
NATUURLIJK VERSPREIDINGSGEBIED	Bolivia, Brazilië, Colombia, Ecuador, Frans-Guyana, Guyana, Peru en Suriname.