herpy

PIJLGIFKIKKERS

Sanguine Poison Frog

Allobates zaparo

Allobates zaparo is a small, diurnal frog of about 3 cm from the rainforest at the foot of the Andes in Ecuador and northern Peru. Adults have a red back, black flanks, a blue belly and yellow spots at the limbs and mimic the poisonous *Ameerega bilinguis* and *Ameerega parvula*, but are not poisonous themselves. The species is only rarely kept, so the care is based on the climate and on its relative *Allobates femoralis*.

Familie: Aromobatidae Vroegere naam: *Epipedobates zaparo; Phyllobates zaparo; Dendrobates zaparo*

KERNGEGEVENS

HET DIER

Volwassen grootte	**3 cm** kop-romplengte
Levensduur	**5-10 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif	**Geen bekend**
Moeilijkheidsgraad	**Ervaren**

VERZORGING

Min. terrarium	**60 × 40 × 40 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**24-27 °C**
Temperatuur 's nachts	**21-23 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Eieren per legsel	**8-20**
Uitkomen	**15-20 dagen**
Larvenperiode	**40-50 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Draagt de kikkervisjes**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	Little is known about the care of *Allobates zaparo*. The species is only rarely kept, and captive breeding is not described.
TEMPERATUUR	Keep the terrarium at 24-27 °C by day and 21-23 °C at night. The figures follow the climate of Ecuador's lowlands at the foot of the Andes. The variation over the year should be small, 1-2 °C. There is no winter rest.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 70-80 % in the middle of the day and close to 100 % in the morning and evening.
WATER EN WATERKWALITEIT	Provide a shallow glass dish with 1-2 cm of water that the frogs can easily climb out of. The dish is used by the adults when the tadpoles are to be deposited.
VERLICHTING EN UVB	The light is first and foremost for the plants: 10-12 hours a day with LED light with a plant-suitable spectrum. It is not documented that *Allobates zaparo* needs UVB.
VOEDING	In the wild ants are an important part of the diet.
GEDRAG EN OMGANG	*Allobates zaparo* is diurnal and lives in the leaf litter. Males call from the top of the leaves to attract females.
NATUURLIJK VERSPREIDINGSGEBIED	Ecuador en Peru.