herpy

PIJLGIFKIKKERS

Pleasing Poison Frog

Ameerega bassleri

Ameerega bassleri is a diurnal poison frog of just under 4 cm from the humid forests at the eastern foot of the Andes in northern Peru. The male guards the eggs and carries the tadpoles to the water, and the species must be kept cooler than most poison frogs. Little is known about its care, and the species is vulnerable in the wild.

Familie: Dendrobatidae Vroegere naam: *Epipedobates bassleri; Dendrobates bassleri; Phobobates bassleri*

KERNGEGEVENS

HET DIER

Volwassen grootte	**4 cm** kop-romplengte
Levensduur	**8-12 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud en bergbos**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**In bladoksels (bijv. bromelia's)**
Geslachtsrijp	**12-13 maanden**
Eieren per legsel	**40**
Uitkomen	**10 dagen**
Larvenperiode	**50-60 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

VERZORGING

Min. terrarium	**50 × 50 × 55 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**18-21 °C**
Temperatuur 's nachts	**13-17 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Insecten en kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Kwetsbaar (VU)**

Verzorging in het kort

HET TERRARIUM	Little is known about the care of *Ameerega bassleri*. The species lives in humid forest by streams at 300-1,200 m, where the air is cooler than for most poison frogs.
TEMPERATUUR	The species must not be kept too warm. By day 18-21 °C suits, and at night the temperature should fall 4-6 °C, to about 13-17 °C. The difference between the seasons is only 1-2 °C.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity must be 70-100 %. Morning and evening it is close to 100 % with mist, while around midday it may fall to about 70 %.
WATER EN WATERKWALITEIT	A water bowl with 1-2 cm of water must be in the vivarium, and a permanent water area or small containers are used for the tadpoles when breeding. Use rainwater or reverse osmosis water and change the water several times a week.
VERLICHTING EN UVB	The species lives on the forest floor, where little light reaches. LED light with a plant-suitable spectrum and a 10-12 hour day length is enough.
VOEDING	The species takes the usual small feeder insects: fruit flies, springtails, newly hatched crickets, the smallest waxworm larvae and sweep-netted small insects from the meadow.
GEDRAG EN OMGANG	The species is diurnal and lives on the ground. The male is territorial and calls to defend his area, so the animals need many hiding places and broken lines of sight.
NATUURLIJK VERSPREIDINGSGEBIED	Peru.