herpy

PIJLGIFKIKKERS

Pleasing Poison Frog

Ameerega bassleri

Ameerega bassleri is a diurnal poison frog of just under 4 cm from the humid forests at the eastern foot of the Andes in northern Peru. The male guards the eggs and carries the tadpoles to the water, and the species must be kept cooler than most poison frogs. Little is known about its care, and the species is vulnerable in the wild.

Familie: Dendrobatidae Vroegere naam: *Epipedobates bassleri; Dendrobates bassleri; Phobobates bassleri*

KERNGEGEVENS

HET DIER

Volwassen grootte	**4 cm** kop-romplengte
Levensduur	**8-12 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud en bergbos**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Gevorderd**

VERZORGING

Min. terrarium	**50 × 50 × 55 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**18-21 °C**
Temperatuur ’s nachts	**13-17 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Insecten en kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**In bladoksels (bijv. bromelia’s)**
Geslachtsrijp	**12-13 maanden**
Eieren per legsel	**40**
Uitkomen	**10 dagen**
Larvenperiode	**50-60 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Kwetsbaar (VU)**

Over de soort

Ameerega bassleri (Melin, 1941) is a medium-sized poison frog in the genus *Ameerega*. Adults measure about 35-38 mm from snout to vent, and the female is slightly larger than the male. There are several local forms, named after the locality, for example 'Saposa' and 'Sisa'. Sex differences other than size and the male's call are not documented.

The species was formerly called *Dendrobates bassleri*, *Phyllobates bassleri*, *Epipedobates bassleri* and *Phobobates bassleri*. The species name honours the American geologist Harvey Bassler. *Ameerega pepperi*, *Ameerega cainarachi* and *Ameerega yoshina* closely resemble the species and are best told apart by the male's call.

Poison frogs in the genus *Ameerega* obtain the alkaloids in their skin from small arthropods in their diet. Captive-bred animals that do not receive these arthropods are not toxic. Wild-caught animals are considered mild: wash your hands before and after contact, and do not touch your eyes or mouth.

Verspreiding en leefgebied

A. bassleri is found only in Peru, in the Amazon drainage at the Río Huallaga, at elevations from about 270 to 1,200 m. The range includes the national parks Cordillera Azul and Cordillera Escalera, among others.

> **Natuurlijk verspreidingsgebied:** Peru.

LEEFGEBIED

The species lives in humid tropical forest in the foothill lowlands and in the mountain forests, in both undisturbed and slightly disturbed forest. The animals are rarely found more than 300 m from a stream. At low elevations they keep to damp places close to brooks, and higher up they are less tied to the water. The tadpoles grow up in small forest pools and brooks. The climate is humid and cooler than in the Amazon lowlands, and the temperature varies only a little through the year.

The species can be very common locally, but is vulnerable because of loss of habitat to agriculture and cattle farming. Collecting for the pet trade is not considered a threat to the species as a whole, but individual colour variants can quickly be overharvested.

Verzorging

HET TERRARIUM

Little is known about the care of *Ameerega bassleri*. The species lives in humid forest by streams at 300-1,200 m, where the air is cooler than for most poison frogs. *Ameerega pepperi* and *Ameerega cainarachi* live in the same part of Peru and have similar requirements. In the genus, *Ameerega trivittata* is the best known in captivity, but it comes from warmer lowlands, so its figures cannot be used directly.

A rainforest vivarium of at least 50 × 50 × 55 cm (length, width, height) suits two adults. Build the floor as a bioactive set-up with a drainage layer, mesh and damp soil, covered by a thick layer of leaf litter. Furnish with bromeliads, moss, branches, roots and cork bark, so that there are many hiding places and the lines of sight are broken. The lid must have mesh so that the air does not stand still.

TEMPERATUUR

The species must not be kept too warm. By day 18-21 °C suits, and at night the temperature should fall 4-6 °C, to about 13-17 °C. The difference between the seasons is only 1-2 °C. Heat above 25 °C should be avoided, so place the vivarium somewhere cool, and avoid heat from the light sources. In warm weather a fan and extra misting can keep the temperature down.

LUCHTVOCHTIGHEID EN SPROEIEN

Humidity must be 70-100 %. Morning and evening it is close to 100 % with mist, while around midday it may fall to about 70 %. Mist or use a rain system and fogger a couple of times a day with rainwater or reverse osmosis water, and let the vivarium dry out a little between mistings so that mould does not form.

Lees meer: Luchtvochtigheid, vernevelen en ventilatie

WATER EN WATERKWALITEIT

A water bowl with 1-2 cm of water must be in the vivarium, and a permanent water area or small containers are used for the tadpoles when breeding. Use rainwater or reverse osmosis water and change the water several times a week. The water must be cool and clean; natural breeding sites are small forest pools and brooks. A filter is not necessary in a small bowl.

VERLICHTING EN UVB

The species lives on the forest floor, where little light reaches. LED light with a plant-suitable spectrum and a 10-12 hour day length is enough. The need for UVB is not documented; a weak UVB bulb (Ferguson zone 1) can be used when there are good hides, so that the animals can withdraw from the light.

Lees meer: Licht en UVB

VOEDING

The species takes the usual small feeder insects: fruit flies, springtails, newly hatched crickets, the smallest waxworm larvae and sweep-netted small insects from the meadow. Feed every other day with as much as the animals eat in about an hour. Dust the feeder insects with calcium with D3 at every second or third feeding and with a vitamin supplement once a week. Young animals get the same feeder insects in smaller sizes.

Lees meer: Calcium, vitaminen en supplementen

GEDRAG EN OMGANG

The species is diurnal and lives on the ground. The male is territorial and calls to defend his area, so the animals need many hiding places and broken lines of sight. A pair can be kept together; several males in the same vivarium cause conflict. Poison frogs should not be handled; move them with a clean glass or a small container when cleaning.

Kweek

GESLACHTSRIJPHEID EN GESLACHTSONDERSCHEID

The animals are sexually mature after 12-13 months. The sexes are best told apart by the male's call; the female is slightly larger. The first clutches from young females are often of poor quality, and the eggs are easily attacked by fungus.

PARING

The male calls to attract the female. During mating the male holds the female around the head. A period of heavier misting that imitates the rain of the forest can set the calling going. After egg-laying the male takes over care of the eggs.

EIEREN EN KIKKERVISJES

In the vivarium the female lays many eggs in bromeliads near the water's edge. A clutch can be over 40 eggs. The male guards the eggs and keeps them moist. When the tadpoles hatch, the male carries them on his back to water. The tadpoles are easy to rear on algae and can also be given prepared tadpole food. They can be reared singly in small containers with cool, soft water that is changed often. The temperature for the tadpoles is not documented; keep them at the same temperature as the adults.

METAMORFOSE

The eggs hatch after about ten days. The tadpoles become small frogs after just under two months, about 55 days. When the forelegs appear, lower the water level, and the metamorphosed froglets need easy access to land and a damp hiding place.

OPFOK VAN DE JONGEN

The newly metamorphosed froglets are small and get springtails and the smallest fruit flies. They are kept cool and damp like the adults, in a vivarium with moss and leaves. They can be kept together in a group as long as there is space. Growth and the time for moving them are not documented.

Gezondheid en veelgemaakte fouten

No disease descriptions exist for the species, so the usual problems in poison frogs apply:

- Too high a temperature, which causes stress and a weakened immune system. The species must be kept cool.
- Fungus on the eggs, especially in the first clutches.
- Chytrid fungus (*Batrachochytrium dendrobatidis*). New animals are quarantined for at least six weeks.
- Lack of calcium and vitamins, which causes weak bones and poor growth.
- Still air and dirty water, which cause bacteria and mould.

Lifespan in captivity is not documented; the figure in the box is an estimate based on related species. Contact a vet with amphibian expertise if an animal stops eating or loses weight.

Lees meer: Quarantaine en hygiëne

Vraag altijd raad aan iemand die er verstand van heeft

Deze gids is een goed beginpunt, maar kan ervaring niet vervangen. Dieren verschillen, en veel van wat er echt toe doet, leer je van mensen die de soort al jaren houden. Praat met een kweker, een ervaren houder of een dierenarts die reptielen en amfibieën kent voordat je het dier haalt - en opnieuw als er iets niet pluis lijkt.

De volledige gids met kaart en updates: herpy.org/en/poison-dart-frogs/ameerega-bassleri