herpy

PIJLGIFKIKKERS

Cainarachi Poison Frog

Ameerega cainarachi

Ameerega cainarachi is a red-backed poison frog of about 3 cm that is known only from the foothills around Tarapoto in northern Peru. The species is threatened in the wild, and little is known about its care, so this guide is based on the climate, the habitat and the close relatives *Ameerega bassleri* and *Ameerega pepperi*.

Familie: Dendrobatidae Vroegere naam: *Epipedobates cainarachi; Epipedobates ardens*

KERNGEGEVENS

HET DIER

Volwassen grootte	**3 cm** kop-romplengte
Levensduur	**10-15 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Geslachtsrijp	**12-13 maanden**
Eieren per legsel	**22-44**
Uitkomen	**10 dagen**
Larvenperiode	**50-60 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

VERZORGING

Min. terrarium	**40 × 40 × 40 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**22-26 °C**
Temperatuur ’s nachts	**19-22 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Insecten en kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Bedreigd (EN)**

Verzorging in het kort

HET TERRARIUM	Little is known about the care of *Ameerega cainarachi*. The species lives in the same area as *Ameerega bassleri* and *Ameerega pepperi*, and the breeding figures are borrowed from them (see Breeding).
TEMPERATUUR	By day 22-26 °C suits, and at night the temperature should fall by up to 5 °C, to about 19-22 °C. The difference between the seasons is only 1-2 °C. There is no winter rest.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity must be 70-100 %: about 70 % around midday and close to 100 % with mist morning and evening. Mist or use a rain system and fogger a couple of times a day.
WATER EN WATERKWALITEIT	A shallow water bowl with 1-2 cm of water is enough, and it must be easy to get in and out of. Use rainwater, reverse osmosis water or dechlorinated tap water, and change the water every second day so that bacteria do not build up.
VERLICHTING EN UVB	The light must first and foremost cover the plants' needs: LED plant-growth light for 10-12 hours a day with a clear night. The need for UVB is not documented; a weak UVB lamp (Ferguson zone 1) is optional.
VOEDING	The frogs take the usual small feeder insects: fruit flies (*Drosophila melanogaster* and *Drosophila hydei*), springtails, small isopods, newly hatched crickets, the smallest waxworm larvae and sweep-netted small insects from the meadow.
GEDRAG EN OMGANG	*Ameerega cainarachi* is diurnal and lives on the forest floor. The males in the genus are territorial and call from fixed spots.
NATUURLIJK VERSPREIDINGSGEBIED	Peru.