herpy

PIJLGIFKIKKERS

Yurimaguas Poison Frog

Ameerega hahneli

Ameerega hahneli is a small, diurnal poison frog of 17-22 mm with pale side stripes and a blue belly, which lives in the rainforest leaf litter across large parts of the Amazon lowlands. The male carries the tadpoles to running water, and the tadpoles need current in the water. The species is common in the wild but rare in captivity.

Familie: Dendrobatidae Vroegere naam: *Epipedobates hahneli; Dendrobates hahneli*

KERNGEGEVENS

HET DIER

Volwassen grootte	**17-22 mm** kop-romplengte
Levensduur	**6-10 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Geslachtsrijp	**7-12 maanden**
Eieren per legsel	**6-33**
Uitkomen	**4-16 dagen**
Larvenperiode	**55-65 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

VERZORGING

Min. terrarium	**50 × 50 × 50 cm** L × B × H voor 4 volwassen dieren
Temperatuur overdag	**23-25 °C**
Temperatuur ’s nachts	**19-21 °C**
Luchtvochtigheid	**65-90%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Insecten en kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM
Only limited knowledge exists about the care of *Ameerega hahneli*, and the requirements may vary between the different forms and possible hidden species.

TEMPERATUUR
Ameerega hahneli needs 23-25 °C by day and 19-21 °C at night. Fluorescent tubes outside the terrarium can supply both light and the necessary warmth. Above 28 °C there is a risk of stress; cool with a fan and extra spraying.

LUCHTVOCHTIGHEID EN SPROEIEN
Humidity must be 65-90 %.

WATER EN WATERKWALITEIT
There must always be a little water to go to, for example in the drainage channel or a shallow water bowl with 1-2 cm of water. In the dry season the channel is not emptied, so the animals still have access to water.

VERLICHTING EN UVB
The light can be fluorescent tubes or LED with a plant-suitable spectrum for 10-12 hours. UVB is not necessary; a weak UVB bulb (Ferguson zone 1) can be used when there are hides so that the animals can withdraw from the light.

VOEDING
The small fruit fly (*Drosophila melanogaster*) is the most important food. Adults also take the large fruit fly (*Drosophila hydei*), but often skip it because it is a little too big.

GEDRAG EN OMGANG
The species is diurnal and lives on the ground, but is most active early in the morning and late in the afternoon and evening; the rest of the day it is shyer.

NATUURLIJK VERSPREIDINGSGEBIED
Brazilië, Bolivia, Peru, Ecuador, Colombia, Venezuela, Guyana, Frans-Guyana en Suriname.