herpy

PIJLGIFKIKKERS

Ruby Poison Frog

Ameerega parvula

Ameerega parvula is a small, diurnal poison frog of 18-24 mm from the rainforest leaf litter in the western Amazon lowlands and at the foot of the Andes. The back is black with dark red spots, and the underside is marbled blue and black. The species is rarely kept, and little is known about its care.

Familie: Dendrobatidae Vroegere naam: *Epipedobates parvulus; Dendrobates parvulus*

KERNGEGEVENS

HET DIER

Volwassen grootte	**18-24 mm** kop-romplengte
Levensduur	**10-15 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Eieren per legsel	**4-16**
Uitkomen	**10-22 dagen**
Larvenperiode	**70-90 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

VERZORGING

Min. terrarium	**40 × 40 × 40 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**25-26 °C**
Temperatuur ‘s nachts	**20-22 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	Little is known about the care of *Ameerega parvula*.
TEMPERATUUR	The daytime temperature is 25-26 °C, and at night it is lowered by 4-5 degrees to 20-22 °C.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity is 70-80 % by day, with a dip to about 70 % around midday, and close to 100 % morning and evening, preferably with mist.
WATER EN WATERKWALITEIT	A shallow water bowl with 1-2 cm of water lets the frogs drink and is necessary if breeding is to take place, because the male deposits the tadpoles there. The bowl must have a slope or stones so that a frog can climb out again.
VERLICHTING EN UVB	The light should first and foremost give the plants a day-night rhythm of about 12 hours, and an LED plant lamp is enough for that. UVB is not necessary for poison frogs, so it is optional.
VOEDING	The food is small, live arthropods: springtails and fruit flies (*Drosophila melanogaster*), micro crickets, sieved meadow plankton and small isopods.
GEDRAG EN OMGANG	The species is diurnal and lives mostly on the ground. The males call and probably defend small territories, as their relatives do, so a pair is the safe way to keep them in a smaller vivarium.
NATUURLIJK VERSPREIDINGSGEBIED	Colombia, Ecuador en Peru.