herpy

PIJLGIFKIKKERS

Ameerega pepperi

Ameerega pepperi is a small, black poison frog with yellow stripes and red, orange or yellow markings from central Peru. It lives by streams in forest at 380-1,000 m and is vulnerable because of clearing. Little is known about its care, which is therefore based on the habitat and close relatives.

Familie: Dendrobatidae

KERNGEGEVENS

HET DIER

Volwassen grootte	**3 cm** kop-romplengte
Levensduur	**8-12 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud en moerasgebied**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Ervaren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Eieren per legsel	**22-44**
Larvenperiode	**35-60 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

VERZORGING

Min. terrarium	**60 × 45 × 45 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**21-25 °C**
Temperatuur 's nachts	**18-21 °C**
Luchtvochtigheid	**75-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Insecten en kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Kwetsbaar (VU)**

Verzorging in het kort

HET TERRARIUM	Little is known about the care of *Ameerega pepperi*. The species lives by streams in humid forest at 380-1,000 m.
TEMPERATUUR	By day the temperature should be 21-25 °C and at night 18-21 °C. It must not rise above 27 °C, as poison frogs tolerate heat poorly. In summer a fan or a cool room may be necessary. There is no winter rest.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity must be 75-100 %. Mist 1-3 times daily with rainwater or reverse osmosis water, or use a rain system, so that the leaf litter becomes damp but not saturated. Excess water must be able to drain away in the drainage layer.
WATER EN WATERKWALITEIT	A shallow water bowl with 1-2 cm of water and a stone or branch as a landing place is sufficient. Use rainwater, reverse osmosis water or dechlorinated water and change it every 2-3 days.
VERLICHTING EN UVB	Provide plant-friendly light for 10-12 hours daily, with shade and hides. UVB is not documented as necessary for the species, but a weak UVB source (Ferguson zone 1) can be used. Take care that the light does not heat the vivarium.
VOEDING	*Ameerega pepperi* is small and takes small feeder insects: fruit flies (mostly *Drosophila melanogaster*, less *Drosophila hydei*), springtails and small isopods.
GEDRAG EN OMGANG	The males are territorial and call from rocks. The species is best kept as a pair in a vivarium with many hides. Frogs should not be handled, because the skin is delicate and handling stresses the animals.
NATUURLIJK VERSPREIDINGSGEBIED	Peru.