herpy

PIJLGIFKIKKERS

Peru Poison Frog

Ameerega petersi

Ameerega petersi is a small, diurnal poison frog of 24-30 mm from the rainforest at the foot of the Andes in eastern Peru and western Brazil. The back is brown with a pale, oblique side stripe. The species is rarely kept, and little is known about its care.

Familie: Dendrobatidae Vroegere naam: *Epipedobates petersi; Dendrobates petersi*

KERNGEGEVENS

HET DIER

Volwassen grootte	**2-3 cm** kop-romplengte
Levensduur	**10-15 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Eieren per legsel	**4-16**
Uitkomen	**10-22 dagen**
Larvenperiode	**70-90 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

VERZORGING

Min. terrarium	**45 × 45 × 45 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**25-28 °C**
Temperatuur ‘s nachts	**21-25 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	Little is known about the care of *Ameerega petersi*, but the species is kept in small numbers, and the basic figures for the vivarium are known. It is kept like other small *Ameerega* species from the rainforest leaf litter.
TEMPERATUUR	The daytime temperature is 25-28 °C, and at night it is lowered by 3-4 degrees to 21-25 °C. The variation over the year is only 1-2 degrees.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity must be 70-100 %: close to 100 % morning and evening, preferably with mist, and about 70-80 % around midday. Mist with reverse osmosis or rainwater morning and evening, or use a fogger or rain system on a timer.
WATER EN WATERKWALITEIT	A shallow water bowl with 1-2 cm of water is enough. The bowl must be shallow enough for a froglet to climb out of it, and have stones or moss at the edge. Change the water often, and clean the bowl with hot water without soap.
VERLICHTING EN UVB	Provide an LED light with a plant-suitable spectrum for 10-12 hours a day. UVB is not necessary for poison frogs; a weak UVB light (Ferguson zone 1) is optional.
VOEDING	The diet consists of small live arthropods from the leaf litter.
GEDRAG EN OMGANG	The males in the genus call and defend small territories. How *Ameerega petersi* behaves in groups has not been described, so a pair is the safest way to keep them.
NATUURLIJK VERSPREIDINGSGEBIED	Peru en Brazilië.