herpy

PIJLGIFKIKKERS

Cordillera Azul Poison Frog

Ameerega rubriventris

Ameerega rubriventris is a small, dark poison dart frog of 21-25 mm with pale side stripes and a red-black or orange-black marbled belly. It lives only in the forests of the Cordillera Azul in central Peru and is threatened. The species is rarely kept, and little is known about its care.

Familie: Dendrobatidae Vroegere naam: *Epipedobates rubriventris*

KERNGEGEVENS

HET DIER

Volwassen grootte	**21-25 mm** kop-romplengte
Levensduur	**10-15 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Eieren per legsel	**4-16**
Uitkomen	**10-22 dagen**
Larvenperiode	**70-90 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

VERZORGING

Min. terrarium	**50 × 50 × 50 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**25-28 °C**
Temperatuur ’s nachts	**21-25 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Bedreigd (EN)**

Verzorging in het kort

HET TERRARIUM	Little is known about the care of *Ameerega rubriventris*.
TEMPERATUUR	The daytime temperature is 25-28 °C, and at night it is lowered by 3-4 degrees to 21-25 °C. The temperature varies only 1-2 degrees over the year, and a winter rest period is not necessary.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 70-80 % by day and close to 100 % morning and evening, ideally with mist. A rain system or a misting machine on a timer makes this easy; otherwise spray thoroughly morning and evening.
WATER EN WATERKWALITEIT	A shallow water bowl with 1-2 cm of water is enough, and it should be easy to get in and out of.
VERLICHTING EN UVB	The light should primarily serve the plants: an LED plant lamp for 10-12 hours a day and a clear night. UVB is not necessary for poison dart frogs; a weak UVB lamp (Ferguson zone 1) is optional.
VOEDING	The frogs eat small live arthropods: fruit flies (*Drosophila melanogaster* and, for adults, also *D. hydei*), springtails, micro crickets, small isopods and sieved meadow plankton.
GEDRAG EN OMGANG	*Ameerega rubriventris* is diurnal and stays mainly on the forest floor, but readily climbs. The male calls with a bird-like peeping, most often 2-3 short whistles in a row.
NATUURLIJK VERSPREIDINGSGEBIED	Peru.