herpy

PIJLGIFKIKKERS

Three-striped Poison Frog

Ameerega trivittata

Ameerega trivittata is a large, diurnal poison frog of 4-6 cm from the forest floor of the Amazon. It is black with two pale green stripes along the flanks, and in a single trip the male often carries 15-30 tadpoles on his back to water. The frog needs a more spacious enclosure than the small poison frogs.

Familie: Dendrobatidae Vroegere naam: *Epipedobates trivittatus; Dendrobates trivittatus; Phobobates trivittatus*

KERNGEGEVENS

HET DIER

Volwassen grootte	**4-6 cm** kop-romplengte
Levensduur	**10-15 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Regenwoud en moerasgebied**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Gevorderd**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Geslachtsrijp	**12-24 maanden**
Eieren per legsel	**15-40**
Uitkomen	**15-22 dagen**
Larvenperiode	**40-90 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Bewaakt de eieren en draagt de kikkervisjes**

VERZORGING

Min. terrarium	**70 × 50 × 50 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**23-25 °C**
Temperatuur ‘s nachts	**19-22 °C**
Luchtvochtigheid	**70-100%**
Water	**Waterbak** diepte 1-2 cm
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Niet bedreigd (LC)**

Verzorging in het kort

HET TERRARIUM	An enclosure of at least 70 × 50 × 50 cm (length, width, height) suits two adult *Ameerega trivittata*.
TEMPERATUUR	Keep the daytime temperature at 23-25 °C, and lower it at night by 3-4 °C to about 19-22 °C. The variation over the year should be small, 1-2 °C.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 80-90 % for most of the day, close to 100 % morning and evening, and it may fall to about 70 % in the middle of the day. A longer time below 70 % is dangerous, especially if the frog has no access to water.
WATER EN WATERKWALITEIT	A shallow water dish with 1-2 cm of water is enough. The dish must be shallow enough for a young frog to climb out, with stones or moss at the edge.
VERLICHTING EN UVB	UVB is not documented as necessary for poison frogs in general, and for this particular species there is no information. A weak UVB light (Ferguson zone 1) is optional.
VOEDING	The diet consists of small live arthropods.
GEDRAG EN OMGANG	The males call loudly and defend small territories, while the females are calm. It is not documented how the species behaves in groups in captivity, so pairs are the safest way to keep them.
NATUURLIJK VERSPREIDINGSGEBIED	Bolivia, Brazilië, Colombia, Guyana, Peru, Suriname en Venezuela.