herpy

PIJLGIFKIKKERS

Cauca Poison Frog

Andinobates bombetes

Andinobates bombetes is a small, diurnal poison dart frog of 17-19 mm from the cloud forests of Colombia's Andes. It is black to dark brown with red or orange markings on the front of the body and depends on water-filled bromeliads. The species is Vulnerable in the wild, and little is known about its care.

Familie: Dendrobatidae Vroegere naam: *Ranitomeya bombetes; Dendrobates bombetes; Minyobates bombetes*

KERNGEGEVENS

HET DIER

Volwassen grootte	**17-19 mm** kop-romplengte
Levensduur	**5-10 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Bergbos**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Ervaren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Eieren per legsel	**1-2**
Uitkomen	**16-24 dagen**
Larvenperiode	**108-112 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Draagt de kikkervisjes**

VERZORGING

Min. terrarium	**45 × 50 × 60 cm** L × B × H voor 5 volwassen dieren
Temperatuur overdag	**22-25 °C**
Temperatuur ’s nachts	**18-21 °C**
Luchtvochtigheid	**80-100%**
Water	**Geen waterbak**
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren of in groepen**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Kwetsbaar (VU)**

Verzorging in het kort

HET TERRARIUM	Little is known about the care of *Andinobates bombetes*. What is known is based on the habitat, a cool and very humid mountain forest with thick leaf litter and bromeliads, and on captive breeding in Colombia.
TEMPERATUUR	The daytime temperature should be 22-25 °C, and at night it may fall to 18-21 °C as in the mountain forest. Keep the temperature below 27 °C.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 80-100 % and never below 70 %. Spray twice a day, morning and afternoon, or use a fogger or rain system with reverse osmosis water or rainwater.
WATER EN WATERKWALITEIT	A water bowl is not needed; the frogs drink droplets from plants and spray water. For the tadpoles, use bromeliads with water in the leaf axils or small plastic cups with 2-3 cm of water.
VERLICHTING EN UVB	The light is primarily for the plants: plant lighting for 10-12 hours a day provides a daily rhythm and growth.
VOEDING	The food is small arthropods: springtails, small fruit flies (*Drosophila melanogaster*), small isopods and newly hatched crickets. Newly metamorphosed froglets take only springtails.
GEDRAG EN OMGANG	*Andinobates bombetes* is diurnal and stays low in the enclosure, in the leaf litter and by the bromeliads. The males call mainly in the morning.
NATUURLIJK VERSPREIDINGSGEBIED	Colombia.