herpy

PIJLGIFKIKKERS

Tolima Poison Frog

Andinobates dorisswansonae

Andinobates dorisswansonae is a small, diurnal poison dart frog of 16-19 mm, black or dark brown with red, orange or yellow spots. It is known only from a small patch of forest in northern Tolima in Colombia. The species is Vulnerable and hardly kept, so its care is based on the cool, wet mountain climate of its habitat.

Familie: Dendrobatidae Vroegere naam: *Ranitomeya dorisswansonae*

KERNGEGEVENS

HET DIER

Volwassen grootte	**16-19 mm** kop-romplengte
Levensduur	**5-10 jaar**
Actief	**Overdag**
Levenswijze	**Bodembewonend**
Leefgebied	**Bergbos**
Huidgif in het wild	**Mild**
Huidgif in gevangenschap	**Geen** Nakweekdieren. Wildvang kan het gif lang behouden
Moeilijkheidsgraad	**Ervaren**

KWEEK

Voortplanting	**Eierleggend**
Ontwikkeling	**Kikkervisjes worden naar het water gedragen**
Legplaats	**Op de grond**
Eieren per legsel	**1-2**
Uitkomen	**16-24 dagen**
Larvenperiode	**108-152 dagen** van uitkomen tot metamorfose
Voeding van de larven	**Algen/planten en kant-en-klaar voer**
Broedzorg	**Draagt de kikkervisjes**

VERZORGING

Min. terrarium	**45 × 45 × 45 cm** L × B × H voor 2 volwassen dieren
Temperatuur overdag	**18-22 °C**
Temperatuur ‘s nachts	**14-17 °C**
Luchtvochtigheid	**85-100%**
Water	**Geen waterbak**
UVB	**Optioneel** Fergusonzone 1
Voeding	**Kleine ongewervelden (springstaarten, fruitvliegen)**
Gehouden	**In paren**

BESCHERMING EN PAPIEREN

CITES	**Bijlage II**
IUCN	**Kwetsbaar (VU)**

Verzorging in het kort

HET TERRARIUM	Little is known about the care of *Andinobates dorisswansonae*, and the species is hardly kept in captivity. The care is based on the habitat: cool, wet mountain forest with a dense canopy, thick leaf litter and bromeliads.
TEMPERATUUR	The temperature has not been measured in the species' habitat. 18-22 °C by day and 14-17 °C at night suit forest at about 1,800 m. The species tolerates heat poorly, and prolonged periods above 25 °C stress the animals.
LUCHTVOCHTIGHEID EN SPROEIEN	Humidity should be 85-100 %. Spray the leaf litter and plants thoroughly once or twice a day with rainwater or reverse osmosis water, or use a fogger on a timer. The substrate should be damp but not waterlogged.
WATER EN WATERKWALITEIT	The species does not need a water section. The animals get moisture from the leaf litter, plants and spraying. The tadpoles grow up in water in the leaf axils of bromeliads.
VERLICHTING EN UVB	Provide 10-12 hours of LED light a day so that the plants thrive. The forest has a dense canopy, so the light should not be strong, and the animals must be able to seek shade.
VOEDING	*Andinobates dorisswansonae* measures only 16-19 mm and eats small arthropods, in the wild especially ants.
GEDRAG EN OMGANG	The species is diurnal and lives on the ground in the leaf litter. Territorial behaviour has not been documented. A pair can be kept together in an enclosure of 45 × 45 × 45 cm.
NATUURLIJK VERSPREIDINGSGEBIED	Colombia.

De volledige gids met kaart en updates: herpy.org/en/poison-dart-frogs/andinobates-dorisswansonae